

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 481 992 A2

(12)

## EUROPEAN PATENT APPLICATION

(43) Date of publication:  
01.12.2004 Bulletin 2004/49

(51) Int Cl.7: C07K 16/18, C12N 15/13,  
C12N 5/10, A61K 39/00,  
A61P 25/28

(21) Application number: 04011466.2

(22) Date of filing: 26.02.2001

(84) Designated Contracting States:  
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE TR  
Designated Extension States:  
AL LT LV MK RO SI

- Bales, Kelly, R.  
Cloverdale IN 46120 (US)
- Paul, Steven, M.  
Carmel IN 46032 (US)
- Tsurushita, Naoya  
Palo Alto CA 94306 (US)
- Vasquez, Maxmillano  
Palo Alto CA 94303 (US)

(30) Priority: 24.02.2000 US 184601 P  
08.12.2000 US 254465 P  
08.12.2000 US 254498 P

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
01913081.4 / 1 257 584

(74) Representative: Grund, Martin, Dr. et al  
Dr. Volker Vossius,  
Patentanwaltskanzlei,  
Geibelstrasse 6  
81679 München (DE)

(71) Applicants:  
• WASHINGTON UNIVERSITY ST. LOUIS  
St. Louis Missouri 63110 (US)  
• Eli Lilly & Company  
Indianapolis, IN 46285 (US)

## Remarks:

This application was filed on 13 - 05 - 2004 as a  
divisional application to the application mentioned  
under INID code 62. The sequence listing, which is  
published as annex to the application documents,  
was filed after the date of filing. The applicant has  
declared that it does not include matter which goes  
beyond the content of the application as filed.

(72) Inventors:  
• Holtzman, David, M.  
St. Louis MO 63141 (US)  
• Demattos, Ronald  
St. Louis MO 63110 (US)

## (54) Humanized antibodies that sequester amyloid beta peptide

(57) A method to treat conditions characterized by  
formation of amyloid plaques both prophylactically and  
therapeutically is described. The method employs hu-  
manized antibodies which sequester soluble A $\beta$  peptide  
from human biological fluids or which preferably specifi-  
cally bind an epitope contained within position 13-28 of  
the amyloid beta peptide A $\beta$ .




Figure 1

**Description**Cross-Reference to Related Applications

5 [0001] This application claims the priority of, United States provisional applications 60/184,601 filed 24 February 2000, 60/254,465, filed 8 December 2000, and 60/254,498, filed 8 December 2000, the contents of each of which are incorporated herein by reference.

Technical Field

10 [0002] The invention relates to humanized antibodies that bind to an epitope between amino acids 13 and 28 of the A $\beta$  peptide and to preventive and therapeutic treatment of conditions associated with beta amyloid, such as Alzheimer's disease, Down's syndrome, and cerebral amyloid angiopathy. More specifically, it concerns use of humanized mon-  
 15 clonal antibodies to sequester amyloid beta (A $\beta$ ) peptide in plasma, brain, and cerebrospinal fluid to prevent accumulation or to reverse deposition of the A $\beta$  peptide within the brain and in the cerebrovasculature and to improve cognition.

Background Art

20 [0003] A number of symptomologies which result in cognitive deficits, stroke, brain hemorrhage, and general mental debilitation appear to be associated with neuritic and cerebrovascular plaques in the brain containing the amyloid beta peptide (A $\beta$ ). Among these conditions are both pre-clinical and clinical Alzheimer's disease, Down's syndrome, and pre-clinical and clinical cerebral amyloid angiopathy (CAA). The amyloid plaques are formed from amyloid beta peptides. These peptides circulate in the blood and in the cerebrospinal fluid (CSF), typically in complexed form with  
 25 lipoproteins. The A $\beta$  peptide in circulating form is composed of 39-43 amino acids (mostly 40 or 42 amino acids) resulting from the cleavage of a common precursor protein, amyloid precursor protein, often designated APP. Some forms of soluble A $\beta$  are themselves neurotoxic and may determine the severity of neurodegeneration and/or cognitive decline (McLean, C. A., *et al.*, *Ann. Neurol.* (1999) 46:860-866; Lambert, M. P., *et al.* (1998) 95:6448-6453; Naslund, J., *J. Am. Med. Assoc.* (2000) 283:1571).

30 [0004] Evidence suggests that A $\beta$  can be transported back and forth between brain and the blood (Gherzi-Egea, J-F., *et al.*, *J. Neurochem.* (1996) 67:880-883; Zlokovic, B. V., *et al.*, *Biochem. Biophys. Res. Comm.* (1993) 67: 1034-1040; Shibata M, *et al.*, *J. Clin. Invest.* (2000) 106:1489-1499). Further A $\beta$  in plaques is in an equilibrium with soluble A $\beta$  in the brain and blood (Kawarabayashi T, *et al.*, *J. Neurosci.* (2001) 21:372-381).

35 [0005] As described in PCT application US00/35681 and U.S. Serial No. 09/153,130 both incorporated herein by reference, total circulating levels of A $\beta$  peptide in CSF are similar in normal individuals and individuals predisposed to exhibit the symptoms of Alzheimer's. However, A $\beta_{42}$  levels are lower on average in individuals with Alzheimer's disease (Nitsch, R. M., *et al.*, *Ann. Neurol.* (1995) 37:512-518). It is known that A $\beta_{42}$  is more prone to aggregate than is A $\beta_{40}$ , and when this happens, adverse consequences such as A $\beta$  deposition in amyloid plaques, conversion of A $\beta$  to toxic soluble forms, nerve cell damage, and behavioral impairment such as dementia ensue (Golde, T.E., *et al.*, *Biochem.*  
 40 *Biophys. Acta.* (2000) 1502:172-187).

[0006] Methods to induce an immune response to reduce amyloid deposits are described in PCT publication WO99/27944 published 10 June 1999. The description postulates that full-length aggregated A $\beta$  peptide would be a useful immunogen. Administration of a A $\beta$  fragment (amino acids 13-28) conjugated to sheep anti-mouse IgG caused no change in cortex amyloid burden, and only one in nine animals that received injections of the A $\beta$  13-28 fragment-  
 45 conjugate showed any lymphoproliferation in response to A $\beta_{40}$ . The application also indicates that antibodies that specifically bind to A $\beta$  peptide could be used as therapeutic agents. However, this appears to be speculation since the supporting data reflect protocols that involve active immunization using, for example, A $\beta_{42}$ . The peptides are supplied using adjuvants and antibody titers formed from the immunization, as well as levels of A $\beta$  peptide and of the precursor peptide, are determined. The publication strongly suggests that A $\beta$  plaque must be reduced in order to alleviate Alzheimer's symptoms, and that cell-mediated processes are required for successful reduction of A $\beta$  plaque.

50 [0007] WO 99/60024, published 25 November 1999, is directed to methods for amyloid removal using anti-amyloid antibodies. The mechanism, however, is stated to utilize the ability of anti-A $\beta$  antibodies to bind to pre-formed amyloid deposits (*i.e.*, plaques) and result in subsequent local microglial clearance of localized plaques. This mechanism was not proved *in vivo*. This publication further states that to be effective against A $\beta$  plaques, anti-A $\beta$  antibodies must gain  
 55 access to the brain parenchyma and cross the blood brain barrier.

[0008] Several PCT applications that relate to attempts to control amyloid plaques were published on 7 December 2000. WO 00/72880 describes significant reduction in plaque in cortex and hippocampus in a transgenic mouse model of Alzheimer' disease when treated using N-terminal fragments of A $\beta$  peptides and antibodies that bind to them, but

not when treated with the A $\beta$  13-28 fragment conjugated to sheep anti-mouse IgG or with an antibody against the 13-28 fragment, antibody 266. The N-terminal directed antibodies were asserted to cross the blood-brain barrier and to induce phagocytosis of amyloid plaques in *in vitro* studies.

[0009] WO 00/72876 has virtually the same disclosure as WO 00/72880 and is directed to immunization with the amyloid fibril components themselves.

[0010] WO 00/77178 describes antibodies that were designed to catalyze the hydrolysis of  $\beta$ -amyloid, including antibodies raised against a mixture of the phenylalanine statine transition compounds Cys-A $\beta$ <sub>10-25</sub>, statine Phe<sub>19</sub>-Phe<sub>20</sub> and Cys-A $\beta$ <sub>10-25</sub> statine Phe<sub>20</sub>-Ala<sub>21</sub> and antibodies raised against A $\beta$ <sub>10-25</sub> having a reduced amide bond between Phe<sub>19</sub> and Phe<sub>20</sub>. This document mentions sequestering of A $\beta$ , but this is speculation because it gives no evidence of such sequestering. Further, the document provides no *in vivo* evidence that administration of antibodies causes efflux of A $\beta$  from the central nervous system, interferes with plaque formation, reduces plaque burden, forms complexes between the antibodies and A $\beta$  in tissue samples, or affects cognition.

[0011] It has been shown that one pathway for A $\beta$  metabolism is via transport from CNS to the plasma (Zlokovic, B. V., *et al.*, *Proc. Natl. Acad. Sci (USA)* (1996) 93:4229-4234; Ghersi-Egea, J-F., *et al.*, *J. Neurochem.* (1996) 67:880-883). Additionally, it has been shown that A $\beta$  in plasma can cross the blood-brain-barrier and enter the brain (Zlokovic, B. V., *et al.*, *Biochem. Biophys. Res. Comm.* (1993) 67:1034-1040). It has also been shown that administration of certain polyclonal and monoclonal A $\beta$  antibodies decreases A $\beta$  deposition in amyloid plaques in the APPV717F transgenic mouse model of Alzheimer's disease (Bard, F., *et al.*, *Nature Med.* (2000) 6:916-919); however, this was said to be due to certain anti-A $\beta$  antibodies crossing the blood-brain-barrier stimulating phagocytosis of amyloid plaques by microglial cells. In Bard's experiments, assays of brain slices *ex vivo* showed that the presence of added A $\beta$  antibody, along with exogenously added microglia, induced phagocytosis of A $\beta$ , resulting in removal of A $\beta$  deposits.

[0012] The levels of both soluble A $\beta$ <sub>40</sub> and A $\beta$ <sub>42</sub> in CSF and blood can readily be detected using standardized assays using antibodies directed against epitopes along the A $\beta$  chain. Such assays have been reported, for example, in U.S. patents 5,766,846; 5,837,672; and 5,593,846. These patents describe the production of murine monoclonal antibodies to the central domain of the A $\beta$  peptide, and these were reported to have epitopes around and including positions 16 and 17. Antibodies directed against the N-terminal region were described as well. Several monoclonal antibodies were asserted to immunoreact with positions 13-28 of the A $\beta$  peptide; these did not bind to a peptide representing positions 17-28, thus, according to the cited patents, establishing that it is this region, including positions 16-17 (the  $\alpha$ -secretase site) that was the target of these antibodies. Among antibodies known to bind between amino acids 13 and 28 of A $\beta$  are mouse antibodies 266, 4G8, and 1C2.

[0013] We have now unexpectedly found that administration of the 266 antibody very quickly and almost completely restores cognition (object memory) in 24-month old hemizygous transgenic mice (APPV717F). Yet, the antibody does not have the properties that the art teaches are required for an antibody to be effective in treating Alzheimer's disease, Down's syndrome, and other conditions related to the A $\beta$  peptide. To our further surprise, we observed that antibodies that bind A $\beta$  between positions 13 and 28 (266 and 4G8) are capable of sequestering soluble forms of A $\beta$  from their bound, circulating forms in the blood, and that peripheral administration of antibody 266 results in rapid efflux of relatively large quantities of A $\beta$  peptide from the CNS into the plasma. This results in altered clearance of soluble A $\beta$ , prevention of plaque formation, and, most surprisingly, improvement in cognition, even without necessarily reducing A $\beta$  amyloid plaque burden, crossing the blood brain barrier to any significant extent, decorating plaque, activating cellular mechanisms, or binding with great affinity to aggregated A $\beta$ .

#### Disclosure of the Invention

[0014] The invention provides humanized antibodies, or fragments thereof, that positively affect cognition in diseases and conditions where A $\beta$  may be involved, such as clinical or pre-clinical Alzheimer's disease, Down's syndrome, and clinical or pre-clinical cerebral amyloid angiopathy. The antibodies or fragments thereof need not cross the blood-brain barrier, decorate amyloid plaque, activate cellular responses, or even necessarily reduce amyloid plaque burden. In another aspect, this invention provides humanized antibodies and fragments thereof that sequester A $\beta$  peptide from its bound, circulating form in blood, and alter clearance of soluble and bound forms of A $\beta$  in central nervous system and plasma. In another aspect, this invention provides humanized antibodies and fragments thereof, wherein the humanized antibodies specifically bind to an epitope between amino acids 13 and 28 of the A $\beta$  molecule. In another aspect, the invention provides humanized antibodies and fragments thereof, wherein the CDR are derived from mouse monoclonal antibody 266 and wherein the antibodies retain approximately the binding properties of the mouse antibody and have *in vitro* and *in vivo* properties functionally equivalent to the mouse antibody (sequences SEQ ID NO:1 through SEQ ID NO:6). In another aspect, this invention provides humanized antibodies and fragments thereof, wherein the variable regions have sequences comprising the CDR from mouse antibody 266 and specific human framework sequences (sequences SEQ ID NO:7 - SEQ ID NO:10), wherein the antibodies retain approximately the binding properties of the mouse antibody and have *in vitro* and *in vivo* properties functionally equivalent to the mouse antibody 266. In

another aspect, this invention provides humanized antibodies and fragments thereof, wherein the light chain is SEQ ID NO:11 and the heavy chain is SEQ ID NO:12.

**[0015]** Also part of the invention are polynucleotide sequences that encode the humanized antibodies or fragments thereof disclosed above, vectors comprising the polynucleotide sequences encoding the humanized antibodies or fragments thereof, host cells transformed with the vectors or incorporating the polynucleotides that express the humanized antibodies or fragments thereof, pharmaceutical formulations of the humanized antibodies and fragments thereof disclosed herein, and methods of making and using the same.

**[0016]** Such humanized antibodies and fragments thereof are useful for sequestering A $\beta$  in humans; for treating and preventing diseases and conditions characterized by A $\beta$  plaques or A $\beta$  toxicity in the brain, such as Alzheimer's disease, Down's syndrome, and cerebral amyloid angiopathy in humans; for diagnosing these diseases in humans; and for determining whether a human subject will respond to treatment using human antibodies against A $\beta$ .

**[0017]** Administration of an appropriate humanized antibody *in vivo* to sequester A $\beta$  peptide circulating in biological fluids is useful for preventive and therapeutic treatment of conditions associated with the formation of A $\beta$ -containing diffuse, neuritic, and cerebrovascular plaques in the brain. The humanized antibody, including an immunologically reactive fragment thereof, results in removal of the A $\beta$  peptide from macromolecular complexes which would normally be relevant in transporting it in body fluids to and from sites where plaques can form or where it can be toxic. In addition, sequestering of plasma A $\beta$  peptide with the antibody or fragment thereof behaves as a "sink," effectively sequestering soluble A $\beta$  peptide in the plasma compartment, and inducing A $\beta$  to enter the plasma from locations in the central nervous system (CNS). By sequestering A $\beta$  in the blood, net efflux from the brain is enhanced and soluble A $\beta$  is prevented from depositing in insoluble plaques and from forming toxic soluble species in the brain. In addition, insoluble A $\beta$  in plaques which is in equilibrium with soluble A $\beta$  can be removed from the brain through a sequestering effect in the blood. Sequestering the A $\beta$  peptide with the antibody also enhances its removal from the body and inhibits toxic effects of soluble A $\beta$  in the brain and the development and further accumulation of insoluble A $\beta$  as amyloid in plaques. The antibodies useful in the invention do not cross the blood-brain barrier in large amounts ( $\leq 0.1\%$  plasma levels). In addition, humanized antibodies used in the invention, when administered peripherally, do not need to elicit a cellular immune response in brain when bound to A $\beta$  peptide or when freely circulating to have their beneficial effects. Further, when administered peripherally they do not need to appreciably bind aggregated A $\beta$  peptide in the brain to have their beneficial effects.

**[0018]** Thus, in one aspect, the invention is directed to a method to treat and to prevent conditions characterized by the formation of plaques containing beta-amyloid protein in humans, which method comprises administering, preferably peripherally, to a human in need of such treatment a therapeutically or prophylactically effective amount of humanized monoclonal antibody or immunologically reactive fragment thereof, which antibody specifically binds to the mid-region of the A $\beta$  peptide. In another aspect, the invention is directed to a method to inhibit the formation of amyloid plaques and to clear amyloid plaques in humans, which method comprises administering to a human subject in need of such inhibition an effective amount of a humanized antibody that sequesters A $\beta$  peptide from its circulating form in blood and induces efflux out of the brain as well as altered A $\beta$  clearance in plasma and the brain. In additional aspects, the invention is directed to such humanized antibodies, including immunologically effective portions thereof, and to methods for their preparation.

**[0019]** The invention also includes methods of reversing cognitive decline, improving cognition, treating cognitive decline, and preventing cognitive decline in a subject diagnosed with clinical or pre-clinical Alzheimer's disease, Down's syndrome, or clinical or pre-clinical cerebral amyloid angiopathy, comprising administering to the subject an effective amount of a humanized antibody of the invention.

**[0020]** The invention also includes use of a humanized antibody of the invention for the manufacture of a medicament, including prolonged expression of recombinant sequences of the antibody or antibody fragment in human tissues, for treating, preventing, or reversing Alzheimer's disease, Down's syndrome, or cerebral amyloid angiopathy; for treating, preventing, or reversing cognitive decline in clinical or pre-clinical Alzheimer's disease, Down's syndrome, or clinical or pre-clinical cerebral amyloid angiopathy; or to inhibit the formation of amyloid plaques or the effects of toxic soluble A $\beta$  species in humans.

**[0021]** The invention is related to the surprising observation that within a short period of time after administration of an antibody of the present invention, relatively large quantities of A $\beta$  efflux from the central nervous system to the blood. Thus, this invention includes methods to assess the response of a human subject to treatment with an antibody that binds A $\beta$  or a fragment thereof, comprising: a) administering the antibody or a fragment thereof to the subject; and b) measuring the concentration of A $\beta$  in the subject's blood.

**[0022]** The invention also includes a method of treating a human subject with an antibody that binds A $\beta$  or a fragment thereof, comprising: a) administering a first amount of the antibody or fragment thereof to the subject; b) within 3 hours to two weeks after administering the first dose, measuring the concentration of A $\beta$  in the subject's blood; c) if necessary, calculating a second amount of antibody or fragment thereof based on the result of step b), which second amount is the same as or different than the first amount; and d) administering the second amount of the antibody or fragment.

[0023] The invention also includes a method of assessing in a human subject the efficacy of an antibody that binds to A $\beta$ , or a fragment thereof, for inhibiting or preventing A $\beta$  amyloid plaque formation, for reducing A $\beta$  amyloid plaque, for reducing the effects of toxic soluble A $\beta$  species, or for treating a condition or a disease associated with A $\beta$  plaque, comprising: a) obtaining a first sample of the subject's plasma or CSF; b) measuring a baseline concentration of A $\beta$  in the first sample; c) administering the antibody or fragment thereof to the subject; d) within 3 hours to two weeks after administering the antibody or fragment thereof, obtaining a second sample of the subject's plasma or CSF; and e) measuring the concentration of A $\beta$  in the second sample; wherein, efficacy is related to the quantity of A $\beta$  bound to the antibody in the blood and the concentration of A $\beta$  in the CSF.

#### Brief Description of the Drawings

#### [0024]

Figure 1 shows the percentage of the A $\beta$  peptide withdrawn from human cerebrospinal fluid through a dialysis membrane by Mab 266 as a function of the molecular weight cutoff of the dialysis membrane.

Figure 2 shows the concentration of A $\beta_{\text{Total}}$  found in the plasma of an APPV717F transgenic mouse after injection with either 200  $\mu$ g or 600  $\mu$ g of Mab 266 as a function of time.

Figure 3A shows the quantity of A $\beta$  peptide deposition in the cortex in APPV717F transgenic mice treated with saline, mouse IgG, or Mab 266. Figure 3B shows correlation of these results with parental origin.

Figure 4 shows the polynucleotide sequences for expressing humanized 266 light chain from plasmid pVk-Hu266 and the single amino acid codes for the expressed humanized 266 light chain (corresponding to SEQ ID NO:11 when mature).

Figure 5 shows the polynucleotide sequences for expressing humanized 266 heavy chain from plasmid pVgl-Hu266 and the single amino acid codes for the expressed humanized 266 heavy chain (corresponding to SEQ ID NO:12 when mature).

Figure 6 is a plasmid map of pVk-Hu266.

Figure 7 is a plasmid map of pVgl-Hu266.

#### Modes of Carrying Out the Invention

[0025] The A $\beta$  peptides that circulate in human biological fluids represent the carboxy terminal region of a precursor protein encoded on chromosome 21. It has been reported from the results of *in vitro* experiments that the A $\beta$  peptide has poor solubility in physiological solutions, since it contains a stretch of hydrophobic amino acids which are a part of the region that anchors its longer precursor to the lipid membranes of cells. It is thus not surprising that circulating A $\beta$  peptide is normally complexed with other moieties that prevent it from aggregating. This has resulted in difficulties in detecting circulating A $\beta$  peptide in biological fluids.

[0026] The above-mentioned patent documents (U.S. patents 5,766,846; 5,837,672 and 5,593,846) describe the preparation of antibodies, including a monoclonal antibody, designated clone 266 which was raised against, and has been shown to bind specifically to, a peptide comprising amino acids 13-28 of the A $\beta$  peptide. The present applicants have found that antibodies that bind within this region, in contrast to antibodies that bind elsewhere in the amino acid sequence of A $\beta$ , are able to sequester the soluble A $\beta$  peptide very effectively from macromolecular complexes. This sequestration will effect net A $\beta$  peptide efflux from the CNS, alter its clearance in CNS and plasma, and reduce its availability for plaque formation. Thus, antibodies of this specificity, modified to reduce their immunogenicity by converting them to a humanized form, offer the opportunity to treat, both prophylactically and therapeutically, conditions that are associated with formation of beta-amyloid plaques. These conditions include, as noted above, pre-clinical and clinical Alzheimer's, Down's syndrome, and pre-clinical and clinical cerebral amyloid angiopathy.

[0027] As used herein, the word "treat" includes therapeutic treatment, where a condition to be treated is already known to be present and prophylaxis - *i.e.*, prevention of, or amelioration of, the possible future onset of a condition.

[0028] By "monoclonal antibodies that bind to the mid-region of A $\beta$  peptide" is meant monoclonal antibodies (Mab or Mabs) that bind an amino acid sequence representing an epitope contained between positions 13-28 of A $\beta$ . The entire region need not be targeted. As long as the antibody binds at least an epitope within this region (especially, *e.g.*, including the  $\alpha$ -secretase site 16-17 or the site at which antibody 266 binds), such antibodies are effective in the method of the invention.

[0029] By "antibody" is meant a monoclonal antibody *per se*, or an immunologically effective fragment thereof, such as an F<sub>ab</sub>, F<sub>ab'</sub>, or F<sub>(ab')<sub>2</sub></sub> fragment thereof. In some contexts, herein, fragments will be mentioned specifically for emphasis; nevertheless, it will be understood that regardless of whether fragments are specified, the term "antibody" includes such fragments as well as single-chain forms. As long as the protein retains the ability specifically to bind its intended target, and in this case, to sequester A $\beta$  peptide from its carrier proteins in blood, it is included within the term

"antibody." Also included within the definition "antibody" for example, are single chain forms, generally designated F<sub>v</sub> regions, of antibodies with this specificity. Preferably, but not necessarily, the antibodies useful in the invention are produced recombinantly, as manipulation of the typically murine or other non-human antibodies with the appropriate specificity is required in order to convert them to humanized form. Antibodies may or may not be glycosylated, though glycosylated antibodies are preferred. Antibodies are properly cross-linked via disulfide bonds, as is well-known.

[0030] The basic antibody structural unit is known to comprise a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function.

[0031] Light chains are classified as gamma, mu, alpha, and lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, and define the antibody's isotype as IgG, IgM, IgA, IgD and IgE, respectively. Within light and heavy chains, the variable and constant regions are joined by a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 10 more amino acids.

[0032] The variable regions of each light/heavy chain pair form the antibody binding site. Thus, an intact antibody has two binding sites. The chains all exhibit the same general structure of relatively conserved framework regions (FR) joined by three hypervariable regions, also called complementarily determining regions or CDRs. The CDRs from the two chains of each pair are aligned by the framework regions, enabling binding to a specific epitope. From N- terminal to C-terminal, both light and heavy chains comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is in accordance with well known conventions [Kabat "Sequences of Proteins of Immunological Interest" National Institutes of Health, Bethesda, Md., 1987 and 1991; Chothia, et al., *J. Mol. Biol.* 196:901-917 (1987); Chothia, et al., *Nature* 342:878-883 (1989)].

[0033] As is well understood in the art, monoclonal antibodies can readily be generated with appropriate specificity by standard techniques of immunization of mammals, forming hybridomas from the antibody-producing cells of said mammals or otherwise immortalizing them, and culturing the hybridomas or immortalized cells to assess them for the appropriate specificity. In the present case such antibodies could be generated by immunizing a human, rabbit, rat or mouse, for example, with a peptide representing an epitope encompassing the 13-28 region of the A $\beta$  peptide or an appropriate subregion thereof. Materials for recombinant manipulation can be obtained by retrieving the nucleotide sequences encoding the desired antibody from the hybridoma or other cell that produces it. These nucleotide sequences can then be manipulated to provide them in humanized form.

[0034] By "humanized antibody" is meant an antibody that is composed partially or fully of amino acid sequences derived from a human antibody germline by altering the sequence of an antibody having non-human complementarity determining regions (CDR). The simplest such alteration may consist simply of substituting the constant region of a human antibody for the murine constant region, thus resulting in a human/murine chimera which may have sufficiently low immunogenicity to be acceptable for pharmaceutical use. Preferably, however, the variable region of the antibody and even the CDR is also humanized by techniques that are by now well known in the art. The framework regions of the variable regions are substituted by the corresponding human framework regions leaving the non-human CDR substantially intact, or even replacing the CDR with sequences derived from a human genome. Fully human antibodies are produced in genetically modified mice whose immune systems have been altered to correspond to human immune systems. As mentioned above, it is sufficient for use in the methods of the invention, to employ an immunologically specific fragment of the antibody, including fragments representing single chain forms.

[0035] A humanized antibody again refers to an antibody comprising a human framework, at least one CDR from a non-human antibody, and in which any constant region present is substantially identical to a human immunoglobulin constant region, i.e., at least about 85-90%, preferably at least 95% identical. Hence, all parts of a humanized antibody, except possibly the CDRs, are substantially identical to corresponding parts of one or more native human immunoglobulin sequences. For example, a humanized immunoglobulin would typically not encompass a chimeric mouse variable region/human constant region antibody.

[0036] Humanized antibodies have at least three potential advantages over non-human and chimeric antibodies for use in human therapy:

1) because the effector portion is human, it may interact better with the other parts of the human immune system (e.g., destroy the target cells more efficiently by complement-dependent cytotoxicity (CDC) or antibody-dependent cellular cytotoxicity (ADCC)).

2) The human immune system should not recognize the framework or C region of the humanized antibody as foreign, and therefore the antibody response against such an injected antibody should be less than against a totally foreign non-human antibody or a partially foreign chimeric antibody.

3) Injected non-human antibodies have been reported to have a half-life in the human circulation much shorter than the half-life of human antibodies. Injected humanized antibodies will have a half-life essentially identical to

EP 1 481 992 A2

naturally occurring human antibodies, allowing smaller and less frequent doses to be given.

[0037] The design of humanized immunoglobulins may be carried out as follows. When an amino acid falls under the following category, the framework amino acid of a human immunoglobulin to be used (acceptor immunoglobulin) is replaced by a framework amino acid from a CDR-providing non-human immunoglobulin (donor immunoglobulin):

(a) the amino acid in the human framework region of the acceptor immunoglobulin is unusual for human immunoglobulin at that position, whereas the corresponding amino acid in the donor immunoglobulin is typical for human immunoglobulin at that position;

(b) the position of the amino acid is immediately adjacent to one of the CDRs; or

(c) any side chain atom of a framework amino acid is within about 5-6 angstroms (center-to-center) of any atom of a CDR amino acid in a three dimensional immunoglobulin model [Queen, *et al.*, *op. cit.*, and Co, *et al.*, *Proc. Natl. Acad. Sci. USA* 88, 2869 (1991)]. When each of the amino acid in the human framework region of the acceptor immunoglobulin and a corresponding amino acid in the donor immunoglobulin is unusual for human immunoglobulin at that position, such an amino acid is replaced by an amino acid typical for human immunoglobulin at that position.

[0038] A preferred humanized antibody is a humanized form of mouse antibody 266. The CDRs of humanized 266 have the following amino acid sequences:

light chain CDR1:

```

      1           5           10           15
Arg Ser Ser Gln Ser Leu Ile Tyr Ser Asp Gly Asn Ala Tyr Leu His
(SEQ ID NO:1)
```

light chain CDR2:

```

      1           5
Lys Val Ser Asn Arg Phe Ser (SEQ ID NO:2)
```

light chain CDR3:

```

      1           5
Ser Gln Ser Thr His Val Pro Trp Thr (SEQ ID NO:3)
```

heavy chain CDR1:

```

      1           5
Arg Tyr Ser Met Ser (SEQ ID NO:4)
```

heavy chain CDR2

```

      1           5           10           15
Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr Pro Asp Thr Val Lys Gly (SEQ
ID NO:5)
```

and, heavy chain CDR3:

```

      1
Gly Asp Tyr (SEQ ID NO:6)
```

[0039] A preferred light chain variable region of a humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline Vk segments DPK18 and J segment

# EP 1 481 992 A2

Jk1, with several amino acid substitutions to the consensus amino acids in the same human V subgroup to reduce potential immunogenicity:

```

5      1      5      10      15
      Asp Xaa Val Met Thr Gln Xaa Pro Leu Ser Leu Pro Val Xaa Xaa

      20      25      30
      Gly Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Xaa

10     35      40      45
      Tyr Ser Asp Gly Asn Ala Tyr Leu His Trp Phe Leu Gln Lys Pro

      50      55      60
      Gly Gln Ser Pro Xaa Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe

15     65      70      75
      Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp

      80      85      90
      Phe Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Xaa Gly Val

20     95     100     105
      Tyr Tyr Cys Ser Gln Ser Thr His Val Pro Trp Thr Phe Gly Xaa

      110
      Gly Thr Xaa Xaa Glu Ile Lys Arg (SEQ ID NO:7)

25

```

wherein:

```

30     Xaa at position 2 is Val or Ile;
      Xaa at position 7 is Ser or Thr;
      Xaa at position 14 is Thr or Ser;
      Xaa at position 15 is Leu or Pro;
      Xaa at position 30 is Ile or Val;
      Xaa at position 50 is Arg, Gln, or Lys;
35     Xaa at position 88 is Val or Leu;
      Xaa at position 105 is Gln or Gly;
      Xaa at position 108 is Lys or Arg; and
      Xaa at position 109 is Val or Leu.

```

40 [0040] A preferred heavy chain variable region of a humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline VH segments DP53 and J segment JH4, with several amino acid substitutions to the consensus amino acids in the same human subgroup to reduce potential immunogenicity:

45

50

55



EP 1 481 992 A2

1 5 10 15  
 Xaa Val Gln Leu Val Glu Xaa Gly Gly Gly Leu Val Gln Pro Gly  
 5 20 25 30  
 Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser  
 35 40 45  
 Arg Tyr Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu  
 10 50 55 60  
 Xaa Leu Val Ala Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr  
 65 70 75  
 Pro Asp Xaa Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Xaa  
 15 80 85 90  
 Xaa Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Xaa Asp  
 95 100 105  
 Thr Ala Val Tyr Tyr Cys Ala Ser Gly Asp Tyr Trp Gly Gln Gly  
 20 110  
 Thr Xaa Val Thr Val Ser Ser (SEQ ID NO:8)

wherein:

- 25 Xaa at position 1 is Glu or Gln;  
 Xaa at position 7 is Ser or Leu;  
 Xaa at position 46 is Glu, Val, Asp, or Ser;  
 Xaa at position 63 is Thr or Ser;  
 30 Xaa at position 75 is Ala, Ser, Val, or Thr;  
 Xaa at position 76 is Lys or Arg;  
 Xaa at position 89 is Glu or Asp; and  
 Xaa at position 107 is Leu or Thr.

35 [0041] A particularly preferred light chain variable region of a humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline Vk segments DPK18 and J segment Jkl, with several amino acid substitutions to the consensus amino acids in the same human V subgroup to reduce potential immunogenicity:

40

45

50

55

EP 1 481 992 A2

1 5 10 15  
Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu  
5 20 25 30  
Gly Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Ile  
35 40 45  
Tyr Ser Asp Gly Asn Ala Tyr Leu His Trp Phe Leu Gln Lys Pro  
10 50 55 60  
Gly Gln Ser Pro Arg Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe  
65 70 75  
Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp  
15 80 85 90  
Phe Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Val Gly Val  
95 100 105  
Tyr Tyr Cys Ser Gln Ser Thr His Val Pro Trp Thr Phe Gly Gln  
20 110  
Gly Thr Lys Val Glu Ile Lys Arg (SEQ ID NO:9).

[0042] A particularly preferred heavy chain variable region of a humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline VH segments DP53 and J segment JH4:

1 5 10 15  
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly  
30 20 25 30  
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser  
35 35 40 45  
Arg Tyr Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu  
50 55 60  
Glu Leu Val Ala Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr  
40 65 70 75  
Pro Asp Thr Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala  
80 85 90  
Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp  
45 95 100 105  
Thr Ala Val Tyr Tyr Cys Ala Ser Gly Asp Tyr Trp Gly Gln Gly  
50 110  
Thr Leu Val Thr Val Ser Ser (SEQ ID NO:10).

[0043] A preferred light chain for a humanized antibody of the present invention has the amino acid sequence:

EP 1 481 992 A2

	1	5	10	15
	Asp	Val	Val	Met
	Thr	Gln	Ser	Pro
	Leu	Ser	Leu	Pro
	Val	Thr	Leu	
5		20	25	30
	Gly	Gln	Pro	Ala
	Ser	Ile	Ser	Cys
	Arg	Ser	Ser	Gln
	Ser	Leu	Ile	
	35	40	45	
	Tyr	Ser	Asp	Gly
	Asn	Ala	Tyr	Leu
	His	Trp	Phe	Leu
	Gln	Lys	Pro	
10		50	55	60
	Gly	Gln	Ser	Pro
	Arg	Leu	Leu	Ile
	Tyr	Lys	Val	Ser
	Asn	Arg	Phe	
	65	70	75	
	Ser	Gly	Val	Pro
	Asp	Arg	Phe	Ser
	Gly	Ser	Gly	Ser
	Gly	Thr	Asp	
15		80	85	90
	Phe	Thr	Leu	Lys
	Ile	Ser	Arg	Val
	Glu	Ala	Glu	Asp
	Val	Gly	Val	
	95	100	105	
	Tyr	Tyr	Cys	Ser
	Gln	Ser	Thr	His
	Val	Pro	Trp	Thr
	Phe	Gly	Gln	
20		110	115	120
	Gly	Thr	Lys	Val
	Glu	Ile	Lys	Arg
	Thr	Val	Ala	Ala
	Pro	Ser	Val	
	125	130	135	
	Phe	Ile	Phe	Pro
	Pro	Ser	Asp	Glu
	Gln	Leu	Lys	Ser
	Gly	Thr	Ala	
25		140	145	150
	Ser	Val	Val	Cys
	Leu	Leu	Asn	Asn
	Phe	Tyr	Pro	Arg
	Glu	Ala	Lys	
	155	160	165	
	Val	Gln	Trp	Lys
	Val	Asp	Asn	Ala
	Leu	Gln	Ser	Gly
	Asn	Ser	Gln	
30		170	175	180
	Glu	Ser	Val	Thr
	Glu	Gln	Asp	Ser
	Lys	Asp	Ser	Thr
	Tyr	Ser	Leu	
	185	190	195	
	Ser	Ser	Thr	Leu
	Thr	Leu	Ser	Lys
	Ala	Asp	Tyr	Glu
	Lys	His	Lys	
35		200	205	210
	Val	Tyr	Ala	Cys
	Glu	Val	Thr	His
	Gln	Gly	Leu	Ser
	Ser	Pro	Val	
40		215		
	Thr	Lys	Ser	Phe
	Asn	Arg	Gly	Glu
	Cys			

(SEQ ID NO:11).

[0044] A preferred heavy chain for a humanized antibody of the present invention has the amino acid sequence:

	1	5	10	15
	Glu	Val	Gln	Leu
	Val	Glu	Ser	Gly
	Gly	Gly	Leu	Val
	Gln	Pro	Gly	

50

55

## EP 1 481 992 A2

		20		25		30
		Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser				
5		35		40		45
		Arg Tyr Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu				
		50		55		60
		Glu Leu Val Ala Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr				
10		65		70		75
		Pro Asp Thr Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala				
		80		85		90
		Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp				
15		95		100		105
		Thr Ala Val Tyr Tyr Cys Ala Ser Gly Asp Tyr Trp Gly Gln Gly				
		110		115		120
		Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val				
20		125		130		135
		Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala				
		140		145		150
		Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr				
25		155		160		165
		Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe				
		170		175		180
		Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val				
30		185		190		195
		Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys				
		200		205		210
		Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val				
35		215		220		225
		Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro				
		230		235		240
		Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro				
40		245		250		255
		Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr				
		260		265		270
		Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe				
45		275		280		285
		Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys				
		290		295		300
		Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val				
50		305		310		315
		Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys				
55						

5  
 10  
 15  
 20  
 25

		320		325		330
	Cys	Lys	Val	Ser	Asn	Lys
					Ala	Leu
					Pro	Ala
					Pro	Ile
						Glu
						Lys
						Thr
		335		340		345
	Ile	Ser	Lys	Ala	Lys	Gly
					Gln	Pro
					Arg	Glu
					Pro	Gln
						Val
						Tyr
						Thr
		350		355		360
	Leu	Pro	Pro	Ser	Arg	Asp
					Glu	Leu
					Thr	Lys
					Asn	Gln
						Val
						Ser
						Leu
		365		370		375
	Thr	Cys	Leu	Val	Lys	Gly
					Phe	Tyr
					Pro	Ser
					Asp	Ile
						Ala
						Val
						Glu
		380		385		390
	Trp	Glu	Ser	Asn	Gly	Gln
					Pro	Glu
					Asn	Asn
					Tyr	Lys
						Thr
						Thr
						Pro
		395		400		405
	Pro	Val	Leu	Asp	Ser	Asp
					Gly	Ser
					Phe	Phe
					Leu	Tyr
						Ser
						Lys
						Leu
		410		415		420
	Thr	Val	Asp	Lys	Ser	Arg
					Trp	Gln
					Gln	Gly
					Asn	Val
						Phe
						Ser
						Cys
		425		430		435
	Ser	Val	Met	His	Glu	Ala
					Leu	His
					Asn	His
					Tyr	Thr
						Gln
						Lys
						Ser
		440				
	Leu	Ser	Leu	Ser	Pro	Gly
					Lys	

(SEQ ID NO:12).

[0045] Other sequences are possible for the light and heavy chains for the humanized antibodies of the present invention and for humanized 266. The immunoglobulins can have two pairs of light chain/heavy chain complexes, at least one chain comprising one or more mouse complementarity determining regions functionally joined to human framework region segments.

[0046] In another aspect, the present invention is directed to recombinant polynucleotides encoding antibodies which, when expressed, comprise the heavy and light chain CDRs from an antibody of the present invention. As to the human framework region, a framework or variable region amino acid sequence of a CDR-providing non-human immunoglobulin is compared with corresponding sequences in a human immunoglobulin variable region sequence collection, and a sequence having a high percentage of identical amino acids is selected. Exemplary polynucleotides, which on expression code for the polypeptide chains comprising the heavy and light chain CDRs of monoclonal antibody 266 are given in Figures 4 and 5. Due to codon degeneracy and non-critical amino-acid substitutions, other polynucleotide sequences can be readily substituted for those sequences. Particularly preferred polynucleotides of the present invention encode antibodies, which when expressed, comprise the CDRs of SEQ ID NO:1 - SEQ ID NO:6, or any of the variable regions of SEQ ID NO:7 - SEQ ID NO:10, or the light and heavy chains of SEQ ID NO:11 and SEQ ID NO:12.

[0047] The polynucleotides will typically further include an expression control polynucleotide sequence operably linked to the humanized immunoglobulin encoding sequences, including naturally-associated or heterologous promoter regions. Preferably, the expression control sequences will be eukaryotic promoter systems in vectors capable of transforming or transfecting eukaryotic host cells, but control sequences for prokaryotic hosts may also be used. Once the vector has been incorporated into the appropriate host cell line, the host cell is propagated under conditions suitable for high level expression of the nucleotide sequences, and, as desired, the collection and purification of the light chains, heavy chains, light/heavy chain dimers or intact antibodies, binding fragments or other immunoglobulin forms may follow.

[0048] The nucleotide sequences of the present invention capable of ultimately expressing the desired humanized antibodies can be formed from a variety of different polynucleotides (genomic or cDNA, RNA, synthetic oligonucleotides, etc.) and components (e.g., V, J, D, and C regions), as well as by a variety of different techniques. Joining appropriate genomic and synthetic sequences is a common method of production, but cDNA sequences may also be utilized.

[0049] Human constant region DNA sequences can be isolated in accordance with well known procedures from a variety of human cells, but preferably from immortalized B-cells. The CDRs for producing the immunoglobulins of the present invention will be similarly derived from non-human monoclonal antibodies capable of binding to an epitope between amino acids 13 and 28 of the A $\beta$  peptide, which monoclonal antibodies are produced in any convenient mammalian source, including, mice, rats, rabbits, or other vertebrates capable of producing antibodies by well known methods, as described above. Suitable source cells for the polynucleotide sequences and host cells for immunoglobulin

expression and secretion can be obtained from a number of sources well-known in the art.

[0050] In addition to the humanized immunoglobulins specifically described herein, other "substantially homologous" modified immunoglobulins can be readily designed and manufactured utilizing various recombinant DNA techniques well known to those skilled in the art. For example, the framework regions can vary from the native sequences at the primary structure level by several amino acid substitutions, terminal and intermediate additions and deletions, and the like. Moreover, a variety of different human framework regions may be used singly or in combination as a basis for the humanized immunoglobulins of the present invention. In general, modifications of the genes may be readily accomplished by a variety of well-known techniques, such as site-directed mutagenesis.

[0051] Alternatively, polypeptide fragments comprising only a portion of the primary antibody structure may be produced, which fragments possess one or more immunoglobulin activities (e.g., complement fixation activity). These polypeptide fragments may be produced by proteolytic cleavage of intact antibodies by methods well known in the art, or by inserting stop codons at the desired locations in vectors using site-directed mutagenesis, such as after CH1 to produce Fab fragments or after the hinge region to produce F(ab')<sub>2</sub> fragments. Single chain antibodies may be produced by joining VL and VH with a DNA linker.

[0052] As stated previously, the encoding nucleotide sequences will be expressed in hosts after the sequences have been operably linked to (i.e., positioned to ensure the functioning of) an expression control sequence. These expression vectors are typically replicable in the host organisms either as episomes or as an integral part of the host chromosomal DNA. Commonly, expression vectors will contain selection markers, e.g., tetracycline or neomycin, to permit detection of those cells transformed with the desired DNA sequences. *E. coli* is a prokaryotic host useful particularly for cloning the polynucleotides of the present invention. Other microbial hosts suitable for use include bacilli, such as *Bacillus subtilis*, and other enterobacteriaceae, such as *Salmonella*, *Serratia*, and various *Pseudomonas* species. In these prokaryotic hosts, one can also make expression vectors, which will typically contain expression control sequences compatible with the host cell (e.g., an origin of replication). In addition, any of a number of well-known promoters may be present, such as the lactose promoter system, a tryptophan (*trp*) promoter system, a beta-lactamase promoter system, or a promoter system from phage lambda. The promoters will typically control expression, optionally with an operator sequence, and have ribosome binding site sequences and the like, for initiating and completing transcription and translation.

[0053] Other microbes, such as yeast, may also be used for expression. *Saccharomyces* is a preferred host, with suitable vectors having expression control sequences, such as promoters, including 3-phosphoglycerate kinase or other glycolytic enzymes, and an origin of replication, termination sequences and the like as desired.

[0054] In addition to microorganisms, mammalian tissue cell culture may also be used to express and produce the polypeptides of the present invention. Eukaryotic cells are actually preferred, because a number of suitable host cell lines capable of secreting intact immunoglobulins have been developed in the art, and include the CHO cell lines, various COS cell lines, Syrian Hamster Ovary cell lines, HeLa cells, preferably myeloma cell lines, transformed B-cells, human embryonic kidney cell lines, or hybridomas. Expression vectors for these cells can include expression control sequences, such as an origin of replication, a promoter, an enhancer, and necessary processing information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences. Preferred expression control sequences are promoters derived from immunoglobulin genes, SV40, Adenovirus, Bovine Papilloma Virus, cytomegalovirus and the like.

[0055] The vectors containing the nucleotide sequences of interest (e.g., the heavy and light chain encoding sequences and expression control sequences) can be transferred into the host cell by well-known methods, which vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment or electroporation may be used for other cellular hosts.

[0056] Once expressed, the whole antibodies, their dimers, individual light and heavy chains, or other immunoglobulin forms of the present invention can be purified according to standard procedures of the art, including ammonium sulfate precipitation, ion exchange, affinity, reverse phase, hydrophobic interaction column chromatography, gel electrophoresis and the like. Substantially pure immunoglobulins of at least about 90 to 95% homogeneity are preferred, and 98 to 99% or more homogeneity most preferred, for pharmaceutical uses. Once purified, partially or to homogeneity as desired, the polypeptides may then be used therapeutically or prophylactically, as directed herein.

[0057] The antibodies (including immunologically reactive fragments) are administered to a subject at risk for or exhibiting A $\beta$ -related symptoms or pathology such as clinical or pre-clinical Alzheimer's disease, Down's syndrome, or clinical or pre-clinical amyloid angiopathy, using standard administration techniques, preferably peripherally (i.e. not by administration into the central nervous system) by intravenous, intraperitoneal, subcutaneous, pulmonary, transdermal, intramuscular, intranasal, buccal, sublingual, or suppository administration. Although the antibodies may be administered directly into the ventricular system, spinal fluid, or brain parenchyma, and techniques for addressing these locations are well known in the art, it is not necessary to utilize these more difficult procedures. The antibodies of the invention are effective when administered by the more simple techniques that rely on the peripheral circulation system. The advantages of the present invention include the ability of the antibody exert its beneficial effects even though not

provided directly to the central nervous system itself. Indeed, it has been demonstrated herein that the amount of antibody which crosses the blood-brain barrier is <0.1% of plasma levels and that the antibodies of the invention exert their ability to sequester A $\beta$  in the peripheral circulation as well as to alter CNS and plasma soluble A $\beta$  clearance.

[0058] The pharmaceutical compositions for administration are designed to be appropriate for the selected mode of administration, and pharmaceutically acceptable excipients such as dispersing agents, buffers, surfactants, preservatives, solubilizing agents, isotonicity agents, stabilizing agents and the like are used as appropriate. Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton PA, latest edition, incorporated herein by reference, provides a compendium of formulation techniques as are generally known to practitioners. It may be particularly useful to alter the solubility characteristics of the antibodies of the invention, making them more lipophilic, for example, by encapsulating them in liposomes or by blocking polar groups.

[0059] Peripheral systemic delivery by intravenous or intraperitoneal or subcutaneous injection is preferred. Suitable vehicles for such injections are straightforward. In addition, however, administration may also be effected through the mucosal membranes by means of nasal aerosols or suppositories. Suitable formulations for such modes of administration are well known and typically include surfactants that facilitate cross-membrane transfer. Such surfactants are often derived from steroids or are cationic lipids, such as N-[1-(2,3-dioleoyl)propyl-N,N,N-trimethylammoniumchloride (DOTMA) or various compounds such as cholesterol hemisuccinate, phosphatidyl glycerols and the like.

[0060] The concentration of the humanized antibody in formulations from as low as about 0.1% to as much as 15 or 20% by weight and will be selected primarily based on fluid volumes, viscosities, and so forth, in accordance with the particular mode of administration selected. Thus, a typical pharmaceutical composition for injection could be made up to contain 1 mL sterile buffered water of phosphate buffered saline and 1-100 mg of the humanized antibody of the present invention. The formulation could be sterile filtered after making the formulation, or otherwise made microbiologically acceptable. A typical composition for intravenous infusion could have a volume as much as 250 mL of fluid, such as sterile Ringer's solution, and 1-100 mg per mL, or more in antibody concentration. Therapeutic agents of the invention can be frozen or lyophilized for storage and reconstituted in a suitable sterile carrier prior to use. Lyophilization and reconstitution can lead to varying degrees of antibody activity loss (e.g. with conventional immune globulins, IgM antibodies tend to have greater activity loss than IgG antibodies). Dosages may have to be adjusted to compensate. The pH of the formulation will be selected to balance antibody stability (chemical and physical) and comfort to the patient when administered. Generally, pH between 4 and 8 is tolerated.

[0061] Although the foregoing methods appear the most convenient and most appropriate for administration of proteins such as humanized antibodies, by suitable adaptation, other techniques for administration, such as transdermal administration and oral administration may be employed provided proper formulation is designed.

[0062] In addition, it may be desirable to employ controlled release formulations using biodegradable films and matrices, or osmotic mini-pumps, or delivery systems based on dextran beads, alginate, or collagen.

[0063] In summary, formulations are available for administering the antibodies of the invention and are well-known in the art and may be chosen from a variety of options.

[0064] Typical dosage levels can be optimized using standard clinical techniques and will be dependent on the mode of administration and the condition of the patient.

[0065] The following examples are intended to illustrate but not to limit the invention.

[0066] The examples hereinbelow employ, among others, a murine monoclonal antibody designated "266" which was originally prepared by immunization with a peptide composed of residues 13-28 of human A $\beta$  peptide. The antibody was confirmed to immunoreact with this peptide, but had previously been reported to not react with the peptide containing only residues 17-28 of human A $\beta$  peptide, or at any other epitopes within the A $\beta$  peptide. The preparation of this antibody is described in U.S. patent 5,766,846, incorporated herein by reference. As the examples here describe experiments conducted in murine systems, the use of murine monoclonal antibodies is satisfactory. However, in the treatment methods of the invention intended for human use, humanized forms of the antibodies with the immunospecificity corresponding to that of antibody 266 are preferred.

#### Example 1

##### Sequestration of Added A $\beta$ Peptide in Human Fluids

[0067] Samples of human cerebrospinal fluid (CSF) (50  $\mu$ l) and human plasma (50  $\mu$ l) were incubated for 1 hour at room temperature as follows:

1. alone;
2. along with 5 ng A $\beta$  40 peptide; or
3. 5 ng A $\beta$  40 peptide plus 1 mg monoclonal antibody 266 (described, for example, in U.S. patent 5,766,846 incorporated herein by reference).

[0068] The samples were then electrophoresed on a 4-25% non-denaturing gradient gel, *i. e.*, non-denaturing gradient electrophoresis (NDGGE) and transferred to nitrocellulose. The blots were then stained with Ponceau S or, for Western blot, probed with biotin-labeled monoclonal antibody (3D6) which is directed against the first five amino acids of A $\beta$  peptide, developed with streptavidin-horse radish peroxidase and detected by enhanced chemiluminescence (ECL). The hydrated diameters of the materials contained in bands on the blots were estimated using Pharmacia molecular weight markers. Thus, if the A $\beta$  peptide is bound to other molecules, it would run at the size of the resulting complex.

[0069] Western blots of CSF either with or without 5 ng A $\beta$  peptide shows no evidence of the A $\beta$  peptide in response to detection mediated by antibody 3D6. Similar results are obtained for human plasma. This was true despite the fact that A $\beta$  peptide could be detected by SDS-PAGE followed by Western blot using the same technique and on the same CSF samples. Presumably, the detection of A $\beta$  peptide was prevented by interactions between this peptide and other factors in the fluids tested. However, when Mab 266 is added to the incubation, characteristic bands representing sequestered A $\beta$  peptide complexed to the antibody are present both in plasma and in CSF. The major band is at approximately 11 nm hydrated diameter, corresponding to antibody monomer with an additional smaller band at 13 nm corresponding to antibody dimer.

## Example 2

### Specificity of the Sequestering Antibody

[0070] Samples containing 50  $\mu$ l of human CSF or 10  $\mu$ l of APPV717F CSF were used. APPV717F are transgenic mice representing a mouse model of Alzheimer's disease in which the human amyloid precursor protein transgene with a familial Alzheimer's disease mutation is expressed and results in the production of human A $\beta$  peptide in the central nervous system.

[0071] The samples were incubated with or without various Mabs (1  $\mu$ g) for 1 hour at room temperature and then electrophoresed on a 4-25% NDGGE and blotted onto nitrocellulose as described in Example 1. The antibodies were as follows:

Mab 266 (binds to positions 13-28);  
 Mab 4G8 (binds to positions 17-24);  
 QCBpan (rabbit polyclonal for positions 1-40);  
 mouse IgG (non-specific);  
 Mab 3D6 (binds to positions 1-5);  
 Mab 21F12 (binds to positions 33-42);  
 Mab 6E10 (binds to positions 1-17); and  
 QCB<sub>40, 42</sub> (rabbit polyclonals to A $\beta$ <sub>40</sub> and A $\beta$ <sub>42</sub>).

[0072] Detection of the A $\beta$  peptide antibody complex was as described in Example 1 - biotin labeled 3D6 (to the A $\beta$  peptide N-terminus) followed by streptavidin-HRP and ECL. Similar detection in human CSF incubated with Mab 266, in some instances substituted QCB<sub>40, 42</sub>, which binds to the carboxyl terminus of A $\beta$  peptide, for 3D6.

[0073] The results showed that of the antibodies tested, only Mab 4G8 and Mab 266 permitted the detection of A $\beta$  peptide.

[0074] The results showed that for human CSF, only Mab 266 and Mab 4G8 were able to sequester in detectable amounts of an antibody A $\beta$  complex (again, without any antibody, no A $\beta$  is detected). Mab 266 was also able to produce similar results to those obtained with human CSF with CSF from APPV717F transgenic mice. A $\beta$  peptide could be sequestered in human CSF using Mab 266 regardless of whether 3D6 or QCB<sub>40, 42</sub> antibody was used to develop the Western blot.

## Example 3

### Demonstration of A $\beta$ Peptide -266 Complex by Two-Dimensional Electrophoresis

[0075] A sample containing 50 ng A $\beta$ <sub>40</sub> peptide was incubated with 2  $\mu$ g Mab 266 at 37°C for 3 hours. A corresponding incubation of Mab 266 alone was used as a control.

[0076] The samples were then subjected to 2-dimensional gel electrophoresis.

[0077] In the first dimension, the incubated samples were subjected to NDGGE as described in Example 1. The polyacrylamide gel was then cut into individual lanes perpendicular to the direction of the first dimensional flow and gel separation under denaturing/reducing conditions by SDS-PAGE (Tricine urea gel) was performed in the second



dimension. The presence of the bands was detected either by Ponceau-S staining (any protein) or by specific development using 6E10 Mab (Senetek, Inc.) and biotinylated anti-mouse A $\beta$  in the HRP-based detection system.

[0078] Ponceau-S staining of the nitrocellulose blots after transfer permitted visualization of the heavy and light chains of Mab 266 alone. It was confirmed that A $\beta$  peptide was in a complex with Mab 266 as a band at 4 kD was observed that aligns with the size of full-length Mab 266 seen after the first dimension NDGGE.

#### Example 4

##### Demonstration of Non-Equivalence of Binding and Sequestration

[0079] A $\beta$  peptide as it circulates in plasma and CSF is thought to be contained in a complex with proteins, including apolipoprotein E. The present example demonstrates that antibodies to apoE, while able to bind to the complex, do not sequester apoE from the remainder of the complex.

[0080] ApoE complexes (500 ng) were incubated with Mab or polyclonal antibodies to apoE (2  $\mu$ g) at 37°C for one hour. The incubated samples were then subjected to NDGGE using the techniques described in Example 1. Following NDGGE, Western blotting was performed with affinity purified goat anti-apoE antibodies with detection by ECL. When no antibody is present, apoE can be detected at 8-13 nm consistent with its presence in lipoprotein particles. The presence of monoclonal or polyclonal antibodies to apoE results in a population shift of apoE to a larger molecular species, a "super shift". This demonstrates that the antibodies to apoE did not sequester, i.e., remove apoE from a lipoprotein particle, rather they bind to apoE on the lipoproteins creating a larger molecular species.

#### Example 5

##### Sequestration of A $\beta$ is Not Perturbed by Anti-apoE Antibodies

[0081] A sample of 100  $\mu$ l human CSF was incubated either with Mab 266 alone, or with polyclonal anti-apoE, or with both antibodies for 60 minutes at 37°C. The samples were then analyzed by NDGGE as described in Example 1 and the detection of bands performed as described in Example 1.

[0082] The results show that as long as Mab 266 was added to the sample, the band at approximately 11 nm diameter characteristic of the sequestered 266-A $\beta$  peptide complex was visible. This is the case whether or not anti-apoE is present. This band, demonstrating sequestered A $\beta$ , also appears if 50 ng of A $\beta$  peptide is added to the incubation mixture in the presence of Mab 266. Thus, alteration of the molecular weight of apoE by the presence of anti-apoE antibodies does not interfere with sequestration of A $\beta$  peptide by Mab 266.

#### Example 6

##### Sequestration of A $\beta$ Peptide *In Vivo*

##### [0083]

A. Transgenic APP<sup>V717F</sup> mice, also termed PDAPP mice, over-express a mutant form of human APP protein. These mice produce human A $\beta$  in the CNS and have elevated levels of human A $\beta$  peptide circulating in the CSF and plasma. Eight month old mice were injected intravenously with saline or 100  $\mu$ g of Mab 266. They were bled 10 minutes after initial injection and again at 20 hours after initial injection.

Samples containing 20  $\mu$ l of plasma from each animal were analyzed by NDGGE and Western blot with antibody 3D6 as described in Example 1. The saline injected animals did not show the presence of the characteristic 11 nm sequestered A $\beta$  peptide band either after 10 minutes or 20 hours. However, the two animals that were injected with Mab 266 did show the appearance of this band after 20 hours.

B. Two month old APP<sup>V717F</sup> mice were used in this study. At day zero, the mice received either no Mab 266, 1 mg Mab 266, or 100  $\mu$ g of this antibody. Plasma samples were taken two days prior to administration of the antibodies and on days 1, 3, 5 and 7. The plasma samples were subjected to NDGGE followed by Western blotting and detection with 3D6 as described in Example 1. At all time points following administration of Mab 266, the 266/A $\beta$  complex was detected unless the plasma sample had been treated with protein G, which binds to immunoglobulin, thus effectively removing the Mab 266. Consistent levels of complex over the time period tested were found except for a slight drop-off at day seven in animals injected with 100  $\mu$ g of Mab 266; in general, the levels in animals administered 100  $\mu$ g were consistently lower than those found in the mice administered 1 mg of this antibody.

C. Two two-month old APP<sup>V717F</sup> mice were administered 1 mg of Mab 266 intravenously and a 25  $\mu$ l plasma sample was taken from each. The plasma sample was subjected to NDGGE followed by Western blot as described above

except that binding with biotinylated 3D6 was followed by detection with streptavidin<sup>125I</sup>(Amersham) and exposure to a phosphorimaging screen. The level of complex was estimated in comparison to a standard curve using known amounts of A $\beta$ <sub>40</sub> complexed with saturating levels of Mab 266 and detected similarly. The amount of A $\beta$  peptide bound to Mab 266 was estimated at approximately 100 ng/ml, representing an increase of approximately 1,000-fold over endogenous A $\beta$  peptide in these mice which had been determined to be about 100 pg/ml. This is also similar to the level of A $\beta$  peptide in APP<sup>V717F</sup> brain prior to A $\beta$  deposition (50-100 ng/g); human APP and human A $\beta$  in APP<sup>V717F</sup> Tg mice are produced almost solely in the brain. Thus, it appears that the presence of Mab 266 in the plasma acts as an A $\beta$  peptide sink facilitating net efflux of A $\beta$  peptide from the CNS into the plasma. This increased net efflux likely results from both increasing A $\beta$  efflux from CNS to plasma and also from preventing A $\beta$  in plasma from re-entering the brain.

The correct size for the sequestered A $\beta$  peptide was confirmed by running 20  $\mu$ L of plasma samples obtained from APP<sup>V717F</sup> mice 24 hours after being injected with 1 mg Mab 266 on TRIS-tricine SDS-PAGE gels followed by Western blotting using anti-A $\beta$  antibody 6E10 prior to, or after, protein G exposure using protein G-bound beads. A band that was depleted by protein G was detected at 4-8 kD, consistent with the presence of monomers and possibly dimers of A $\beta$  peptide.

D. Two month old APP<sup>V717F</sup> mice were treated with either PBS (n=7) or 500  $\mu$ g biotinylated Mab 266 - i.e., m266B (n=9) intraperitoneally. Both prior to and 24 hours after the injection, plasma was analyzed for total A $\beta$  peptide using a modification of the ELISA method of Johnson-Wood, K., et al., *Proc. Natl. Acad. Sci. USA* (1997) 94: 1550-1555; and Bales, K.R., et al., *Nature Genet* (1997) 17:263-264. Total A $\beta$  bound to m266B was measured by using 96-well Optiplates (Packard, Inc.) coated with m3D6. Diluted plasma samples and standards (varying concentrations of A $\beta$ <sub>40</sub> and m266B) were incubated overnight in the coated plates and the amount of total A $\beta$ /m266B complex was determined with the use of <sup>125I</sup>-Streptavidin. In addition, at the 24-hour time point, the plasma samples were first treated with protein G to quantitate A $\beta$  peptide not bound to Mab 266, and A $\beta$ <sub>Total</sub> and A $\beta$ <sub>42</sub> were determined by ELISA in the CSF. In PBS-injected animals, plasma A $\beta$  peptide levels were 140 pg/ml both before and after injection. Plasma levels were similar in the Mab 266-injected mice prior to injection, but levels of A $\beta$  peptide not bound to Mab 266 were undetectable at 24 hours post injection.

**[0084]** Levels in the CSF were also measured, CSF represents an extracellular compartment within the CNS and concentration of molecules in the CSF reflects to some extent the concentration of substances in the extracellular space of the brain. CSF was isolated from the cisterna magna compartment. Mice were anesthetized with pentobarbital and the musculature from the base of the skull to the first vertebrae was removed. CSF was collected by carefully puncturing the arachnoid membrane covering the cistern with a micro needle under a dissecting microscope and withdrawing the CSF into a polypropylene micropipette. At 24 hours post injection, an increase in total A $\beta$  peptide in the CSF of Mab 266-injected mice was found, and an approximately two-fold increase in A $\beta$ <sub>42</sub> as compared to PBS injected mice was obtained in the CSF. This was confirmed using denaturing gel electrophoresis followed by Western blotting with A $\beta$ <sub>42</sub>-specific antibody 21F12.

**[0085]** In an additional experiment, three month old APP<sup>V717F</sup> Tg mice were injected with either PBS or Mab 266 intravenously and both A $\beta$ <sub>40</sub> and A $\beta$ <sub>42</sub> levels were assessed in the CSF as follows:

**[0086]** For measurement of A $\beta$ <sub>40</sub>, the monoclonal antibody m2G3, specific for A $\beta$ <sub>40</sub> was utilized. The ELISA described (Johnson-Wood, K., et al., *Proc. Natl. Acad. Sci. USA* (1997) 94:1550-1555) was modified into an RIA by replacing the Streptavidin-HRP reagent with <sup>125I</sup>-Streptavidin. For plasma and CSF samples, the procedure was performed under non-denaturing conditions that lacked guanidine in the buffers. For assessment of carbonate soluble and insoluble A $\beta$  in brain homogenate, samples were homogenized with 100 mM carbonate, 40 mM NaCl, pH 11.5 (4°C), spun at 10,000 x g for 15 min, and A $\beta$  was assessed in the supernatant (soluble) and the pellet (insoluble) fractions as described (Johnson-Wood, K., et al., *Proc. Natl. Acad. Sci. USA* (1997) 94:1550-1555) and listed above. The measurement of A $\beta$ /Mab 266 complex in plasma was performed by a modified RIA. Mice were injected with biotinylated Mab 266 (Mab 266B) and plasma was isolated at multiple time points. Total A $\beta$  bound to Mab 266 was measured by using 96-well Optiplates (Packard, Inc.) coated with m3D6. Diluted plasma samples and standards (varying concentrations of A $\beta$ <sub>40</sub> and Mab 266B) were incubated overnight in the coated plates and the amount of total A $\beta$ /Mab 266B complex was determined with the use of <sup>125I</sup>-Streptavidin.

**[0087]** Three hours following the intravenous injection of Mab 266, there was a two-fold increase in CSF A $\beta$ <sub>40</sub> levels and a non-significant increase in A $\beta$ <sub>42</sub>. However, at both 24 and 72 hours there was a two to three-fold increase in both A $\beta$ <sub>40</sub> and A $\beta$ <sub>42</sub> in the CSF. Similar results were obtained using denaturing gel analysis followed by A $\beta$  Western blotting of pooled CSF. The efflux of A $\beta$  through brain interstitial fluid, which is reflected to some degree by CSF levels, likely accounts for the observed increase in CSF A $\beta$ .

**[0088]** It is significant that the change in CSF A $\beta$  peptide levels cannot be due to entry of Mab 266 into the CSF since the levels measured 24 hours after injection, which are less than 0.1% plasma levels of Mab 266, are insufficient to account for the changes. These results suggest A $\beta$  peptide is withdrawn from the brain parenchyma into the CSF by

the presence of the antibody in the bloodstream.

[0089] Forms of A $\beta$  peptide which are soluble in PBS or carbonate buffer were measured in cerebral cortical homogenates in the same mice which had been injected with Mab 266 and in which the CSF was analyzed as described above. Similar increases in these soluble forms in the cortical homogenates were observed.

#### Example 7

##### Mab 266 Acts as an A $\beta$ Peptide Sink *In Vitro*

[0090] A dialysis chamber was constructed as an *in vitro* system to test the ability of Mab 266 to act as a sink for A $\beta$  peptide. One mL of human CSF was placed in the top chamber of a polypropylene tube separated by a dialysis membrane with a specified cutoff in the range 10-100 kD from a bottom chamber containing 75  $\mu$ L PBS with or without 1  $\mu$ g of Mab 266.

[0091] It appeared that equilibrium was reached after 3 hours, as determined by subjecting material in the bottom chamber to acid urea gels followed by Western blotting for A $\beta$  peptide with 6E10 at various time points. Samples were denatured in formic acid to a final concentration of 80% (vol/vol) and reduced with  $\beta$ -mercaptoethanol (1%). Samples were electrophoresed (anode to cathode) in a 0.9 M acetic acid running buffer through a 4% to 35% polyacrylamide gradient gel containing 6 M urea, 5% (vol/vol) glacial acetic acid, and 2.5% TEMED. The acidic pH of the gel was neutralized prior to transfer to nitrocellulose. Subsequently, standard Western blotting techniques were used to identify A $\beta$ . The bands detected correspond to 4 kD.

[0092] The amount of A $\beta$  removed from the top chamber was thus determined by ELISA analysis of both top and bottom chambers (n=4) after 3 hours. The results for various molecular weight cutoffs in the presence and absence of Mab 266 are shown in Figure 1. As shown, while only minimal amounts of A $\beta$  peptide crossed the membrane when PBS was placed in the bottom chamber, 50% of the A $\beta$  peptide was sequestered in the bottom chamber when Mab 266 was present and the molecular weight cutoff was 25 kD; increasing amounts crossed as the molecular weight cutoff increased to 100 kD, when almost 100% of the A $\beta$  peptide was drawn across the membrane.

[0093] It was also observed that the anti-N-terminal A $\beta$  antibodies 3D6 and 10D5 were able to draw A $\beta$  peptide across the membrane in this system, though not able to sequester A $\beta$  peptide in the assays described in Example 1. These results show that antibodies to the A $\beta$  peptide have sufficient affinity under these conditions to sequester the peptide in physiological solutions away from other binding proteins, but that Mabs such as 266 which are immunoreactive with an epitope in positions 13-28 are substantially more efficient and bind with higher affinity.

[0094] In similar assays, astrocyte-secreted apoE4 which was purified as described by DeMattos, R.B., *et al.*, *J. Biol. Chem.* (1998) 273:4206-4212; Sun, Y., *et al.*, *J. Neurosci.* (1998) 18:3261-3272, had a small but statistically significant effect in increasing the mass of A $\beta$  peptide in the bottom chamber. No apparent effect was observed when polyclonal IgG or BSA was substituted for Mab 266.

#### Example 8

##### Flux of A $\beta$ Peptide into Plasma from the CNS

##### [0095]

A. One  $\mu$ g of A $\beta$ <sub>40</sub> was dissolved into 5  $\mu$ L of rat CSF to keep it soluble and was then injected into the subarachnoid space of the cisterna magna of wild-type Swiss-Webster mice which had previously received IV injections of either PBS (n=3) or 200  $\mu$ g of biotinylated Mab 266 (n=3). At different time-points following treatment, A $\beta$ <sub>Total</sub> in the plasma of the mice was determined by A $\beta$  ELISA, using 3D6 as the coating antibody and standards of A $\beta$  mixed with an excess of biotinylated 266. Each plasma sample was spiked with an excess of biotinylated 266 after removal from each animal for A $\beta$  detection in the ELISA. In the PBS-injected mice, minimally detectable amounts of the peptide at levels of 0.15 ng/ml were detected as peak values after 30-60 minutes, after which the levels were essentially zero. In the mice administered Mab 266, however, plasma A $\beta$  peptide reached levels 330-fold higher than those detected in PBS-injected mice after 60 minutes (approximately 50 ng/ml) and reached values of approximately 90 ng/ml after 180 minutes.

B. This procedure was repeated using either 200  $\mu$ g (n=3) or 600  $\mu$ g (n=3) injected IV into two-month-old APP<sup>V717F</sup> mice. Mab 266 was injected i.v. into 3 month old APP<sup>V717F</sup> +/- mice with the above doses. Prior to and at different time-points following i.v. injection, the plasma concentration of A $\beta$  bound to Mab 266 was determined by RIA. The detailed results from one illustrative mouse are shown in Figure 2.

[0096] It was found that the concentration of A $\beta$  bound to the monoclonal antibody Mab 266 increased from basal

levels of 150 pg/ml to levels of over 100 ng/ml by four days. By analyzing early time points on the curve, it was determined that the net rate of entry of  $A\beta_{Total}$  into plasma of the APPV717F Tg mice was 42 pg/ml/minute in the presence of saturating levels of the antibody.

[0097] The effects of Mab 266 on plasma  $A\beta$  levels in both wild type and APPV717F Tg mice as well as the effects of the antibody on  $A\beta$  concentration in CSF show that the presence of circulating Mab 266 results in a change in the equilibrium of  $A\beta$  flux or transport between the CNS and plasma.

#### Example 9

##### Mab 266 Effect on $A\beta$ in the Brain

[0098] Four month old APPV717F+/+ mice were treated every 2 weeks for 5 months with IP injections of saline, Mab 266 (500  $\mu$ g), or control mouse IgG (100  $\mu$ g, Pharmigen). The mice were sacrificed at nine months of age, and  $A\beta$  deposition in the cortex was determined. The % area covered by  $A\beta$ -immunoreactivity, as identified with a rabbit pan- $A\beta$  antibody (QCB, Inc.), was quantified in the cortex immediately overlying the dorsal hippocampus as described by Holtzman, D.M., *et al.*, *Ann. Neurol.* (2000) 97:2892-2897. The results are shown in Figure 3A. At this age, about half of each group has still not begun to develop  $A\beta$  deposition. However, the % of mice with >50%  $A\beta$  burden in the cortex was significantly less ( $P=0.02$ , Chi-square test) in the 266-treated group. While APPV717F mice can develop large amounts of  $A\beta$  deposits by nine months, there is great variability with about 50% showing no deposits and about 50% showing substantial deposits. In PBS and IgG treated animals, 6/14 and 5/13 mice had greater than 50% of the cortex covered by  $A\beta$  staining, while only one of 14 mice treated with Mab 266 had this level of staining. Almost 50% of the animals in all groups still had not developed  $A\beta$  deposition by 9 months of age. The latter appears to be due to parental origin of individual mice in our cohort since even though all mice studied were confirmed to be APPV717F+/+, high levels of  $A\beta$  deposition was observed only in mice derived from 4/8 breeding pairs (High pathology litters). Mice derived from the other 4 breeding pairs were virtually free of  $A\beta$  deposits (Low pathology litters). Using parental origin as a covariate, there was a strong, significant effect of m266 in reducing  $A\beta$  deposition ( $p=0.0082$ , Fig. 3B).

#### Example 10

##### Peripherally injected Mab 266 does not bind to plaques in APPV717F Tg mice

[0099] To determine whether Mab 266 injected i.p. over 5 months was bound to  $A\beta$  in brain, brain sections from 9 month old APPV717F+/+ Tg mice which contained  $A\beta$  deposits and had been treated with either Mab 266, saline, or control IgG were utilized. Tissue processing and immunostaining was performed as described (Bales, K.R., *et al.*, *Nature Genet.* (1997) 17:263-264). Tissue from all groups of animals was incubated with fluorescein-labeled anti-mouse IgG (Vector, Inc.) and then examined under a fluorescent microscope. No specific staining of  $A\beta$  deposits was seen in any of the groups. In contrast, when applying Mab 266 to sections prior to incubation of the sections with anti-mouse IgG,  $A\beta$  deposits were clearly detected.

#### Example 11

##### Effect of administration of antibody 266 on cognition in 24-month old transgenic, hemizygous PDAPP mice

[0100] Sixteen hemizygous transgenic mice (APPV717F) were used. The mice were approximately 24 months old at the start of the study. All injections were intraperitoneal (i.p.). Half the mice received weekly injections of phosphate buffered saline (PBS, "Control") and the other half received 500 micrograms of mouse antibody 266 dissolved in PBS. Injections were made over a period of seven weeks (42 days) for a total of six injections. Three days following the last injection, the behavior of the animals was assessed using an object recognition task, essentially as described in J.-C. Dodart, *et al.*, *Behavioral Neuroscience*, 113 (5) 982-990 (1999). A recognition index  $(T_B \times 100)/(T_B - T_A)$  was calculated. Results are shown below in Table 1.

**Table 1. Descriptive statistics for recognition index**

	N	Recognition Index (minutes)		
		Mean	Standard Deviation	Standard Error
Control (PBS)	8	71.2**	8.80	3.11
Antibody 266	8	54.35	7.43	2.62

\*\* p=0.0010

[0101] Administration of 500 micrograms of antibody 266 weekly to 24 month old, hemizygous, transgenic mice was associated with a significant change in behavior. Antibody treated transgenic mice had recognition indices which were similar to wildtype control animals [J.-C. Dodart, *et al*]. The difference in the recognition index was statistically significant at the 0.001 probability level. The increased recognition index is an indication that treatment with an antibody that binds to the beta amyloid peptide in the region of amino acids 13-28 will reverse the behavioral impairments that had been documented in this mouse model of Alzheimer's Disease. Therefore, the administration of antibodies that bind beta amyloid peptide in the region of amino acids 13-28 will treat diseases such as Alzheimer's disease and Down's syndrome and will halt the cognitive decline typically associated with disease progression.

[0102] The amyloid burden (% area covered by immunoreactive material after staining with anti-A $\beta$  antibodies 3D6 or 21F12) was quantified in the cortex immediately overlying the hippocampus including areas of the cingulate and parietal cortex from the brains of the 24 month-old animals treated with mouse antibody 266 for seven weeks, as described above. The results are presented in the table below. The differences between the treatment groups are not statistically significant.

**Table 2. Amyloid plaque burden in APP<sup>V717F</sup> +/- mice following treatment with mouse 266 anti-A $\beta$  antibody**

	N	Plaque Burden (%)			
		Using 3D6		Using 21F12	
		Mean	Standard Error	Mean	Standard Error
Control (PBS)	7	44.3	5.93	0.77	0.14
Antibody 266	8	38.0	2.96	0.93	0.11

[0103] For these very old animals, treatment with mouse antibody 266 did not result in a significantly different amyloid burden compared with the PBS-treated group, measured using either 3D6 or using 21F12. Furthermore, the A $\beta$  burden was substantially greater and significantly increased compared with the amyloid burden in younger animals (see below) who were not able to discriminate a novel object from a familiar one in the object recognition task. Most surprisingly, these results demonstrate that anti-A $\beta$  antibodies can reverse cognitive deficits without the need to reduce amyloid burden *per se*.

[0104] After 7 weeks of treatment, the recognition index of the m266-treated group was not significantly different than what would be expected for a wild type cohort of 24 month old mice! This indicates a complete reversal of cognitive decline in these transgenic animals.

#### Example 12

##### Effect of administration of antibody 266 on cognition in young transgenic, hemizygous PDAPP mice

[0105] Fifty-four (54) homozygous, transgenic mice (APP<sup>V717F</sup>) were used. Twenty-three (23) mice were approximately two months old at the start of the study. The remaining mice were approximately four months old at the start of the study. The duration of treatment was five months. Thus, at study termination, the mice were either approximately

seven (7) months old or approximately nine (9) months old.

[0106] All injections were intraperitoneal (i.p.). Each mouse in "PBS" control groups received a weekly injection of phosphate buffered saline (PBS; 200  $\mu$ L). Each mouse in the "IgG" control groups received a weekly injection of IgG1 $\kappa$ 1 isotype control (100  $\mu$ g/mouse/week). Each mouse in the "High Dose" groups received a weekly injection of 500 microgram of antibody 266 dissolved in PBS ("HD"). Each mouse in the "Low Dose" group received a weekly injection of 100 micrograms of antibody 266 dissolved in PBS ("LD"). Three days following the last injection, the behavior of the animals was assessed using an object recognition task, as described in Example 10 above, and a discrimination index was calculated as the difference between the time spent on a novel object and the time spent on a familiar object. Results are shown below in Table 3. The data are grouped by the age of the mice at the end of the study.

**Table 3. Descriptive statistics for discrimination index**

		Discrimination Index (minutes)		
	N	Mean	Standard Deviation	Standard Error
<b>7 months old</b>				
PBS	7	2.12	4.22	1.59
IgG	8	0.81	3.64	1.29
HD	8	10.04*	6.52	2.30
<b>9 months old</b>				
PBS	7	1.87	3.54	1.34
IgG	8	0.96	3.51	1.24
LD	8	10.75*	6.44	2.28
HD	8	12.06***	7.82	2.76

\*  $p < 0.05$

\*\*\*  $p < 0.0001$

[0107] Taken together these data support the conclusion that administration of antibody 266, an antibody directed against the central domain of A $\beta$ , attenuates plaque deposition in 7-9 month old APP<sup>V717F</sup> transgenic mice, as well as reverses the behavioral impairments previously characterized. Treatment of patients with an antibody directed against the central domain of the A $\beta$  peptide will inhibit or prevent cognitive decline typically associated with disease progression, and will reverse it.

[0108] The discrimination index for treated animals was not significantly different than what would be expected for wild type mice of the same age. Thus, just as in older animals (Example 11), treatment with m266 completely reversed cognitive decline in these younger transgenic animals.

### Example 13

#### Synthesis of Humanized Antibody 266

[0109] **Cells and antibodies.** Mouse myeloma cell line Sp2/0 was obtained from ATCC (Manassas, VA) and maintained in DME medium containing 10% FBS (Cat # SH32661.03, HyClone, Logan, UT) in a 37°C CO<sub>2</sub> incubator. Mouse 266 hybridoma cells were first grown in RPMI-1640 medium containing 10% FBS (HyClone), 10 mM HEPES, 2 mM glutamine, 0.1 mM non-essential amino acids, 1 mM sodium pyruvate, 25  $\mu$ g/ml gentamicin, and then expanded in serum-free media (Hybridoma SFM, Cat # 12045-076, Life Technologies, Rockville, MD) containing 2% low Ig FBS (Cat # 30151.03, HyClone) to a 2.5 liter volume in roller bottles. Mouse monoclonal antibody 266 (Mu266) was purified from the culture supernatant by affinity chromatography using a protein-G Sepharose column. Biotinylated Mu266 was prepared using EZ-Link Sulfo-NHS-LC-LC-Biotin (Cat # 21338ZZ, Pierce, Rockford, IL).

**[0110]** Cloning of variable region cDNAs. Total RNA was extracted from approximately  $10^7$  hybridoma cells using TRIzol reagent (Life Technologies) and poly(A)<sup>+</sup> RNA was isolated with the PolyAtract mRNA Isolation System (Promega, Madison, WI) according to the suppliers' protocols. Double-stranded cDNA was synthesized using the SMART<sup>TM</sup>-RACE cDNA Amplification Kit (Clontech, Palo Alto, CA) following the supplier's protocol. The variable region cDNAs for the light and heavy chains were amplified by polymerase chain reaction (PCR) using 3' primers that anneal respectively to the mouse kappa and gamma chain constant regions, and a 5' universal primer provided in the SMART<sup>TM</sup>-RACE cDNA Amplification Kit. For VL PCR, the 3' primer has the sequence:

5' -TATAGAGCTCAAGCTTGGATGGTGGGAAGATGGATACAGTTGGTGC-3'

[SEQ ID NO: 13]

with residues 17- 46 hybridizing to the mouse Ck region. For VH PCR, the 3' primers have the degenerate sequences:

5' -TATAGAGCTCAAGCTTCCAGTGGATAGACCGATGGGGCTGTCGTTTTGGC-3'   
 A G T   
 T

[SEQ ID NO: 14]

with residues 17 - 50 hybridizing to mouse gamma chain CH1. The VL and VH cDNAs were subcloned into pCR4Blunt-TOPO vector (Invitrogen, Carlsbad, CA) for sequence determination. DNA sequencing was carried out by PCR cycle sequencing reactions with fluorescent dideoxy chain terminators (Applied Biosystems, Foster City, CA) according to the manufacturer's instruction. The sequencing reactions were analyzed on a Model 377 DNA Sequencer (Applied Biosystems).

**[0111]** Construction of humanized 266 (Hu266) variable regions. Humanization of the mouse antibody V regions was carried out as outlined by Queen et al. [*Proc. Natl. Acad. Sci. USA* 86:10029-10033 (1988)]. The human V region framework used as an acceptor for Mu266 CDRs was chosen based on sequence homology. The computer programs ABMOD and ENCAD [Levitt, M., *J. Mol. Biol.* 168:595-620 (1983)] were used to construct a molecular model of the variable regions. Amino acids in the humanized V regions that were predicted to have contact with CDRs were substituted with the corresponding residues of Mu266. This was done at residues 46, 47, 49, and 98 in the heavy chain and at residue 51 in the light chain. The amino acids in the humanized V region that were found to be rare in the same V-region subgroup were changed to the consensus amino acids to eliminate potential immunogenicity. This was done at residues 42 and 44 in the light chain.

**[0112]** The light and heavy chain variable region genes were constructed and amplified using eight overlapping synthetic oligonucleotides ranging in length from approximately 65 to 80 bases [He, X. Y., et al., *J. Immunol.* 160: 029-1035 (1998)]. The oligonucleotides were annealed pairwise and extended with the Klenow fragment of DNA polymerase I, yielding four double-stranded fragments. The resulting fragments were denatured, annealed pairwise, and extended with Klenow, yielding two fragments. These fragments were denatured, annealed pairwise, and extended once again, yielding a full-length gene. The resulting product was amplified by PCR using the Expand High Fidelity PCR System (Roche Molecular Biochemicals, Indianapolis, IN). The PCR-amplified fragments were gel-purified and cloned into pCR4Blunt-TOPO vector. After sequence confirmation, the VL and VH genes were digested with MluI and XbaI, gel-purified, and subcloned respectively into vectors for expression of light and heavy chains to make pVk-Hu266 and pVg1-Hu266 (see Figures 6 and 7, respectively, herein) [Co, M. S., et al., *J. Immunol.* 148:1149-1154 (1992)]. The mature humanized 266 antibody expressed from these plasmids has the light chain of SEQ ID NO:11 and the heavy chain of SEQ ID NO:12.

**[0113]** Stable transfection. Stable transfection into mouse myeloma cell line Sp2/0 was accomplished by electroporation using a Gene Pulser apparatus (BioRad, Hercules, CA) at 360 V and 25  $\mu$ F as described (Co et al., 1992). Before transfection, pVk-Hu266 and pVg1-Hu266 plasmid DNAs were linearized using FspI. Approximately  $10^7$  Sp2/0 cells were transfected with 20  $\mu$ g of pVk-Hu266 and 40  $\mu$ g of pVg1-Hu266. The transfected cells were suspended in DME medium containing 10% FBS and plated into several 96-well plates. After 48 hr, selection media (DME medium containing 10% FBS, HT media supplement, 0.3 mg/ml xanthine and 1  $\mu$ g/ml mycophenolic acid) was applied. Approximately 10 days after the initiation of the selection, culture supernatants were assayed for antibody production by ELISA as shown below. High yielding clones were expanded in DME medium containing 10% FBS and further analyzed for antibody expression. Selected clones were then adapted to growth in Hybridoma SFM.

**[0114]** Measurement of antibody expression by ELISA. Wells of a 96-well ELISA plate (Nunc-Immuno plate, Cat #

439454, NalgeNunc, Naperville, IL) were coated with 100  $\mu$ l of 1  $\mu$ g/ml goat anti-human IgG, Fc $\gamma$  fragment specific, polyclonal antibodies (Cat # 109-005-098, Jackson ImmunoResearch, West Grove, PA) in 0.2 M sodium carbonate-bicarbonate buffer (pH 9.4) overnight at 4°C. After washing with Washing Buffer (PBS containing 0.1% Tween 20), wells were blocked with 400  $\mu$ l of Superblock Blocking Buffer (Cat # 37535, Pierce) for 30 min and then washed with Washing Buffer. Samples containing Hu266 were appropriately diluted in ELISA Buffer (PBS containing 1% BSA and 0.1% Tween 20) and applied to ELISA plates (100  $\mu$ l per well). As a standard, humanized anti-CD33 IgG1 monoclonal antibody HuM195 (Co, *et al.*, 1992, *above*) was used. The ELISA plate was incubated for 2 hr at room temperature and the wells were washed with Wash Buffer. Then, 100  $\mu$ l of 1/1,000-diluted HRP-conjugated goat anti-human kappa polyclonal antibodies (Cat # 1050-05, Southern Biotechnology, Birmingham, AL) in ELISA Buffer was applied to each well. After incubating for 1 hr at room temperature and washing with Wash Buffer, 100  $\mu$ l of ABTS substrate (Cat #s 507602 and 506502, Kirkegaard and Perry Laboratories, Gaithersburg, MD) was added to each well. Color development was stopped by adding 100  $\mu$ l of 2% oxalic acid per well. Absorbance was read at 415 nm using an OPTImax microplate reader (Molecular Devices, Menlo Park, CA).

**[0115]** Purification of Hu266. One of the high Hu266-expressing Sp2/0 stable transfectants (clone 1D9) was adapted to growth in Hybridoma SFM and expanded to 2 liter in roller bottles. Spent culture supernatant was harvested when cell viability reached 10% or below and loaded onto a protein-A Sepharose column. The column was washed with PBS before the antibody was eluted with 0.1 M glycine-HCl (pH 2.5), 0.1 M NaCl. The eluted protein was dialyzed against 3 changes of 2 liter PBS and filtered through a 0.2  $\mu$ m filter prior to storage at 4°C. Antibody concentration was determined by measuring absorbance at 280 nm (1 mg/ml = 1.4 A<sub>280</sub>). SDS-PAGE in Tris-glycine buffer was performed according to standard procedures on a 4-20% gradient gel (Cat # EC6025, Novex, San Diego, CA). Purified humanized 266 antibody is reduced and run on an SDS-PAGE gel. The whole antibody shows two bands of approximate molecular weights 25 kDa and 50 kDa. These results are consistent with the molecular weights of the light chain and heavy chain or heavy chain fragment calculated from their amino acid compositions.

#### **Example 14**

##### *In vitro* binding properties of humanized 266 antibody

**[0116]** The binding efficacy of humanized 266 antibody, synthesized and purified as described above, was compared with the mouse 266 antibody using biotinylated mouse 266 antibody in a comparative ELISA. Wells of a 96-well ELISA plate (Nunc-Immuno plate, Cat # 439454, NalgeNunc) were coated with 100  $\mu$ l of  $\beta$ -amyloid peptide (1-42) conjugated to BSA in 0.2 M sodium carbonate/bicarbonate buffer (pH 9.4) (10  $\mu$ g/mL) overnight at 4°C. The A $\beta$ <sub>1-42</sub>-BSA conjugate was prepared by dissolving 7.5 mg of A $\beta$ <sub>1-42</sub>-Cys<sub>43</sub> (C-terminal cysteine A $\beta$ <sub>1-42</sub>, AnaSpec) in 500  $\mu$ l of dimethylsulfoxide, and then immediately adding 1,500  $\mu$ l of distilled water. Two (2) milligrams of maleimide-activated bovine serum albumin (Pierce) was dissolved in 200  $\mu$ l of distilled water. The two solutions were combined, thoroughly mixed, and allowed to stand at room temperature for two (2) hours. A gel chromatography column was used to separate unreacted peptide from A $\beta$ <sub>1-42</sub>-Cys-BSA conjugate.

**[0117]** After washing the wells with phosphate buffered saline (PBS) containing 0.1% Tween 20 (Washing Buffer) using an ELISA plate washer, the wells were blocked by adding 300  $\mu$ l of SuperBlock reagent (Pierce) per well. After 30 minutes of blocking, the wells were washed with Washing Buffer and excess liquid was removed.

**[0118]** A mixture of biotinylated Mu266 (0.3  $\mu$ g/ml final concentration) and competitor antibody (Mu266 or Hu266; starting at 750  $\mu$ g/ml final concentration and serial 3-fold dilutions) in ELISA Buffer were added in triplicate in a final volume of 100  $\mu$ l per well. As a no-competitor control, 100  $\mu$ l of 0.3  $\mu$ g/ml biotinylated Mu266 was added. As a background control, 100  $\mu$ l of ELISA Buffer was added. The ELISA plate was incubated at room temperature for 90 min. After washing the wells with Washing Buffer, 100  $\mu$ l of 1  $\mu$ g/ml HRP-conjugated streptavidin (Cat # 21124, Pierce) was added to each well. The plate was incubated at room temperature for 30 min and washed with Washing Buffer. For color development, 100  $\mu$ l/well of ABTS Peroxidase Substrate (Kirkegaard & Perry Laboratories) was added. Color development was stopped by adding 100  $\mu$ l/well of 2% oxalic acid. Absorbance was read at 415 nm. The absorbances were plotted against the log of the competitor concentration, curves were fit to the data points (using Prism) and the IC<sub>50</sub> was determined for each antibody using methods well-known in the art.

**[0119]** The mean IC<sub>50</sub> for mouse 266 was 4.7  $\mu$ g/mL (three separate experiments, standard deviation = 1.3  $\mu$ g/mL) and for humanized 266 was 7.5  $\mu$ g/mL (three separate experiments, standard deviation = 1.1  $\mu$ g/mL). A second set of three experiments were carried out, essentially as described above, and the mean IC<sub>50</sub> for mouse 266 was determined to be 3.87  $\mu$ g/mL (SD = 0.12  $\mu$ g/mL) and for human 266, the IC<sub>50</sub> was determined to be 4.0  $\mu$ g/mL (SD = 0.5  $\mu$ g/mL). On the basis of these results, we conclude that humanized 266 has binding properties that are very similar to those of the mouse antibody 266. Therefore, we expect that humanized 266 has very similar *in vitro* and *in vivo* activities compared with mouse 266 and will exhibit in humans the same effects demonstrated with mouse 266 in mice.



Example 15In vitro binding properties of mouse antibodies 266 and 4G8

5 [0120] Antibody affinity ( $KD = Kd / Ka$ ) was determined using a BIAcore biosensor 2000 and data analyzed with BIAevaluation (v. 3.1) software. A capture antibody (rabbit anti-mouse) was coupled via free amine groups to carboxyl groups on flow cell 2 of a biosensor chip (CM5) using N-ethyl-N-dimethylaminopropyl carbodiimide and N-hydroxysuccinimide (EDC/NHS). A non-specific rabbit IgG was coupled to flow cell 1 as a background control. Monoclonal antibodies were captured to yield 300 resonance units (RU). Amyloid-beta 1-40 or 1-42 (Biosource International, Inc.) was then flowed over the chip at decreasing concentrations (1000 to 0.1 times KD). To regenerate the chip, bound anti-A $\beta$  antibody was eluted from the chip using a wash with glycine-HCl (pH 2). A control injection containing no amyloid-beta served as a control for baseline subtraction. Sensorgrams demonstrating association and dissociation phases were analyzed to determine Kd and Ka. Using this method, the affinity of mouse antibody 266 for both A $\beta$ <sub>1-40</sub> and for A $\beta$ <sub>1-42</sub> was found to be 4 pM. The affinity of 4G8 for A $\beta$ <sub>1-40</sub> was 23 nM and for A $\beta$ <sub>1-42</sub> was 24 nM. Despite a 6000-fold difference in affinities for A $\beta$ , both 266 and 4G8, which bind to epitopes between amino acids 13 and 28 of A $\beta$ , effectively sequester A $\beta$  from human CSF. Therefore, the location of the epitope is paramount, rather than binding affinity, in determining the ability of an antibody to sequester A $\beta$  and to provide the beneficial and surprising advantages of the present invention.

20 Example 16Epitope mapping of mouse antibody 266 using Biacore methodology and soluble peptides

25 [0121] The BIAcore is an automated biosensor system for measuring molecular interactions [Karlsson R., et al. J. Immunol. Methods 145:229-240 (1991)]. The advantage of the BIAcore over other binding assays is that binding of the antigen can be measured without having to label or immobilize the antigen (i.e. the antigen maintains a more native conformation). The BIAcore methodology was used to assess the binding of various amyloid-beta peptide fragments to mouse antibody 266, essentially as described above in Example 12, except that all dilutions were made with HEPES buffered saline containing Tween 20, a variety of fragments of A $\beta$  (BioSource International) were injected, and a single concentration of each fragment was injected (440 nM).

30 [0122] Amyloid beta fragments 1-28, 12-28, 17-28 and 16-25 bound to mouse antibody 266, while A $\beta$  fragments 1-20, 10-20, and 22-35 did not bind. Fragments 1-20, 10-20 and 22-35 bound to other MAbs with known epitope specificity for those regions of A $\beta$ . Using this methodology, the binding epitope for the mouse antibody 266 appears to be between amino acids 17 and 25 of A $\beta$ . Since binding usually occurs with at least 3 residues of the epitope present, one could further infer that the epitope is contained within residues 19-23.

Example 17In vitro binding properties of humanized antibody 266

40 [0123] The affinity ( $KD = Kd / Ka$ ) of humanized 266 antibody, synthesized and purified as described above, was determined essentially as described above in Example 15. Using this method, the affinity of humanized 266 for A $\beta$ <sub>1-42</sub> was found to be 4 pM.

45

50

55

SEQUENCE LISTING

5 <110> ELI LILLY AND COMPANY and WASHINGTON UNIVERSITY  
 <120> Humanized Antibodies that Sequester Aβ Peptide  
 <130> 53-33T1  
 10 <150> PCT/US01/06191  
 <151> 2001-02-26  
 <150> 60/184,601  
 <151> 2000-02-24  
 15 <150> 60/254,465  
 <151> 2000-12-08  
 <150> 60/254,498  
 <151> 2000-12-08  
 20 <160> 16  
 <170> PatentIn version 3.1  
 25 <210> 1  
 <211> 16  
 <212> PRT  
 <213> Mus sp.  
 30  
 <400> 1  
 Arg Ser Ser Gln Ser Leu Ile Tyr Ser Asp Gly Asn Ala Tyr Leu His  
 1 5 10 15  
 35  
 <210> 2  
 <211> 7  
 40 <212> PRT  
 <213> Mus sp.  
 45  
 <400> 2  
 Lys Val Ser Asn Arg Phe Ser  
 1 5  
 50 <210> 3  
 <211> 9  
 <212> PRT  
 <213> Mus sp.  
 55

5 <400> 3  
 Ser Gln Ser Thr His Val Pro Trp Thr  
 1 5

10 <210> 4  
 <211> 5  
 <212> PRT  
 <213> Mus sp.

15 <400> 4  
 Arg Tyr Ser Met Ser  
 1 5

20 <210> 5  
 <211> 17  
 <212> PRT  
 <213> Mus sp.

25 <400> 5  
 Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr Pro Asp Thr Val Lys  
 1 5 10 15

30 Gly

35 <210> 6  
 <211> 3  
 <212> PRT  
 <213> Mus sp.

40 <400> 6  
 Gly Asp Tyr  
 1

45 <210> 7  
 <211> 113  
 <212> PRT  
 <213> Artificial sequence

50

55

5 <220>  
 <223> Humanized antibodies  
 <220>  
 <221> MISC\_FEATURE  
 10 <222> (2)..(2)  
 <223> xaa at position 2 is Val or Ile  
 15 <220>  
 <221> MISC\_FEATURE  
 <222> (7)..(7)  
 20 <223> xaa at position 7 is Ser or Thr  
 <220>  
 25 <221> MISC\_FEATURE  
 <222> (14)..(14)  
 <223> xaa at position 14 is Thr or Ser  
 30 <220>  
 <221> MISC\_FEATURE  
 <222> (15)..(15)  
 35 <223> xaa at position 15 is Leu or Pro  
 <220>  
 40 <221> MISC\_FEATURE  
 <222> (30)..(30)  
 <223> xaa at position 30 is Ile or Val  
 45 <220>  
 <221> MISC\_FEATURE  
 50 <222> (50)..(50)  
 <223> xaa at position 50 is Arg, Gln, or Lys  
 55 <220>

<221> MISC\_FEATURE  
 <222> (88)..(88)  
 5 <223> xaa at position 88 is val or Leu  
  
 <220>  
 10 <221> MISC\_FEATURE  
 <222> (105)..(105)  
 <223> xaa at position 105 is Gln or Gly  
 15  
 <220>  
 <221> MISC\_FEATURE  
 20 <222> (108)..(108)  
 <223> xaa at position 108 is Lys or Arg  
  
 <220>  
 <221> MISC\_FEATURE  
 25 <222> (109)..(109)  
 <223> xaa at position 109 is val or Leu  
 30  
 <400> 7  
 35 Asp Xaa Val Met Thr Gln Xaa Pro Leu Ser Leu Pro Val Xaa Xaa Gly  
 1. 5 10 15  
 Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Xaa Tyr Ser  
 20 25 30  
 40 Asp Gly Asn Ala Tyr Leu His Trp Phe Leu Gln Lys Pro Gly Gln Ser  
 35 40 45  
 Pro Xaa Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro  
 45 50 55 60  
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
 65 70 75 80  
 50 Ser Arg val Glu Ala Glu Asp xaa Gly val Tyr Tyr Cys Ser Gln Ser  
 85 90 95  
 Thr His val Pro Trp Thr Phe Gly xaa Gly Thr xaa xaa Glu Ile Lys  
 100 105 110  
 55

EP 1 481 992 A2

Arg

5

<210> 8

<211> 112

<212> PRT

10

<213> Artificial sequence

<220>

15

<223> Humanized antibodies

<220>

<221> MISC\_FEATURE

20

<222> (1)..(1)

<223> Xaa at position 1 is Glu or Gln

25

<220>

<221> MISC\_FEATURE

<222> (7)..(7)

30

<223> Xaa at position 7 is Ser or Leu

<220>

35

<221> MISC\_FEATURE

<222> (46)..(46)

<223> Xaa at position 46 is Glu, Val, Asp or Ser

40

<220>

<221> MISC\_FEATURE

<222> (63)..(63)

45

<223> Xaa at position 63 is Thr or Ser

<220>

50

<221> MISC\_FEATURE

<222> (75)..(75)

<223> Xaa at position 75 is Ala, Ser, Val or Thr

55

<220>  
 <221> MISC\_FEATURE  
 5 <222> (76)..(76)  
 <223> Xaa at position 76 is Lys or Arg  
  
 10 <220>  
 <221> MISC\_FEATURE  
 <222> (89)..(89)  
 15 <223> Xaa at position 89 is Glu or Asp  
  
 <220>  
 20 <221> MISC\_FEATURE  
 <222> (107)..(107)  
 <223> Xaa at position 107 is Leu or Thr  
  
 25 <400> 8  
 Xaa Val Gln Leu Val Glu Xaa Gly Gly Gly Leu Val Gln Pro Gly Gly  
 1 5 10 15  
 30 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Arg Tyr  
 20 25 30  
 Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Xaa Leu Val  
 35 35 40 45  
 Ala Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr Pro Asp Xaa Val  
 50 55 60  
 40 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Xaa Xaa Asn Thr Leu Tyr  
 65 70 75 80  
 Leu Gln Met Asn Ser Leu Arg Ala Xaa Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95  
 45 Ala Ser Gly Asp Tyr Trp Gly Gln Gly Thr Xaa Val Thr Val Ser Ser  
 100 105 110  
 50 <210> 9  
 <211> 113  
 <212> PRT  
 55 <213> Artificial sequence

5 <220>  
 <223> Humanized antibodies  
 <400> 9  
 10 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly  
 1 5 10 15  
 Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Ile Tyr Ser  
 20 25 30  
 15 Asp Gly Asn Ala Tyr Leu His Trp Phe Leu Gln Lys Pro Gly Gln Ser  
 35 40 45  
 Pro Arg Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro  
 50 55 60  
 20 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
 65 70 75 80  
 25 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Ser  
 85 90 95  
 Thr His Val Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
 100 105 110  
 30 Arg  
 35 <210> 10  
 <211> 112  
 <212> PRT  
 <213> Artificial sequence  
 40  
 <220>  
 <223> Humanized antibodies  
 45 <400> 10  
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
 1 5 10 15  
 50 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Arg Tyr  
 20 25 30  
 Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Leu Val  
 35 40 45  
 55



EP 1 481 992 A2

Ala Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr Pro Asp Thr Val  
50 55 60

5 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr  
65 70 75 80

10 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
85 90 95

Ala Ser Gly Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser  
100 105 110

15 <210> 11  
<211> 219  
<212> PRT  
20 <213> Artificial sequence

<220>  
25 <223> Humanized antibodies  
<400> 11

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly  
1 5 10 15

30 Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Ile Tyr Ser  
20 25 30

35 Asp Gly Asn Ala Tyr Leu His Trp Phe Leu Gln Lys Pro Gly Gln Ser  
35 40 45

Pro Arg Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro  
50 55 60

40 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
65 70 75 80

45 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Ser  
85 90 95

Thr His Val Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
100 105 110

50 Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu  
115 120 125

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe  
130 135 140

55

EP 1 481 992 A2

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln  
 145 150 155 160  
 5 ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser  
 165 170 175  
 Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu  
 10 180 185 190  
 Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser  
 195 200 205  
 15 Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys  
 210 215  
 <210> 12  
 20 <211> 442  
 <212> PRT  
 <213> Artificial sequence  
 25 <220>  
 <223> Humanized antibodies  
 <400> 12  
 30 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
 1 5 10 15  
 35 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Arg Tyr  
 20 25 30  
 Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Leu Val  
 35 40 45  
 40 Ala Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr Pro Asp Thr Val  
 50 55 60  
 45 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr  
 65 70 75 80  
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95  
 50 Ala Ser Gly Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser  
 100 105 110  
 55 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys  
 115 120 125

EP 1 481 992 A2

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr  
 130 135 140  
 5 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser  
 145 150 155 160  
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser  
 165 170 175  
 10 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr  
 180 185 190  
 Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys  
 195 200 205  
 15 Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys  
 210 215 220  
 20 Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro  
 225 230 235 240  
 Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys  
 245 250 255  
 25 Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp  
 260 265 270  
 30 Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu  
 275 280 285  
 Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu  
 290 295 300  
 35 His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn  
 305 310 315 320  
 40 Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly  
 325 330 335  
 Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu  
 340 345 350  
 45 Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr  
 355 360 365  
 Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn  
 370 375 380  
 50 Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe  
 385 390 395 400  
 55

EP 1 481 992 A2

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn  
405 410 415

5 Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr  
420 425 430

10 Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
435 440

<210> 13

<211> 46

15 <212> DNA

<213> Synthetic nucleotides

20 <400> 13  
tatagagctc aagcttggat ggtgggaaga tggatacagt tgggtgc 46

<210> 14

25 <211> 50

<212> DNA

<213> Synthetic nucleotides

30 <400> 14  
tatagagctc aagcttccag tggatagach gatggggstg tygttttggc 50

35 <210> 15

<211> 16

<212> PRT

<213> Artificial sequence

<400> 15

40 Arg Ser Ser Gln Ser Leu Val Tyr Ser Asp Gly Asn Ala Tyr Leu His  
1 5 10 15

<210> 16

<211> 17

45 <212> PRT

<213> Artificial sequence

<400> 16

50 Gln Ile Asn Ser Val Gly Asn Ser Thr Tyr Tyr Pro Asp Ser Val Lys  
1 5 10 15

Gly

55

Claims

1. A humanized antibody or fragment thereof that specifically binds an epitope contained within positions 13-28 of A $\beta$ .
- 5 2. A humanized antibody or fragment thereof that sequesters A $\beta$  from its bound, circulating form in blood, and alters clearance of soluble and bound forms of A $\beta$  in central the nervous system and plasma.
3. The humanized antibody or fragment thereof of claims 1 or 2 that specifically binds an epitope contained within positions 13-20 of A $\beta$ .
- 10 4. The humanized antibody or fragment thereof of claims 1 or 2 that specifically binds an epitope having an amino acid between positions 17-25 of A $\beta$ .
5. The humanized antibody or fragment thereof of claim 4 that specifically binds an epitope having an amino acid between positions 19-23 of A $\beta$ .
- 15 6. The humanized antibody or fragment thereof of claim 1 or 2 that specifically binds to the epitope of A $\beta$  peptide to which antibody Mab 266 specifically binds.
- 20 7. The humanized antibody or fragment thereof of claim 6, which is humanized Mab 266 or a fragment thereof.
8. The humanized antibody or fragment thereof of claims 4 or 5, wherein the epitope position is determined by the BIAcore method.
- 25 9. A pharmaceutical composition that comprises the humanized antibody or fragment thereof of any one of claims 1-8, and a pharmaceutically acceptable excipient.
10. The use of the humanized antibody or fragment thereof of any one of claims 1-8 for the manufacture of a medication for inhibiting the formation of amyloid plaques or the effects of toxic soluble A $\beta$  species; reducing amyloid plaques or the effects of toxic soluble A $\beta$  species; preventing, treating or reversing cognitive decline; or improving cognition in a human subject.
- 30 11. The use of the humanized antibody or fragment thereof of any one of claims 1-8 for the manufacture of a medication for preventing or treating clinical or pre-clinical Alzheimer's disease, Down's syndrome, or clinical or pre-clinical cerebral amyloid angiopathy in a human subject.
- 35 12. The use of any one of claims 10-11, wherein said antibody or fragment, when administered peripherally to a human subject, need not cross the blood-brain barrier to exert its beneficial effects.
- 40 13. The use of any one of claims 10-11, wherein said antibody or fragment, when administered peripherally to a human subject, does not require cellular responses in the subject's brain to exert its beneficial effects.




Figure 1

**A $\beta$  Levels in Plasma following Intravenous  
266 Injection in APP<sup>V717F</sup> Mice**




Figure 2




Figure 3



Figure 4. pVk-Hu266 polynucleotide sequences for expressing humanized 266 light chain and single amino acid codes for expressed humanized 266 light chain

619 ACGCGTCCACCATGAAGTTGCCCTGTTAGGCTGTTGGTGTCTGATGTTCTGGATTCTGCTCCAGGTGTGATGTTGTGATG  
M K L P V R L L V L M F H I P A S R C D V V M

699 ACCCAGAGCCCACTCTCCCTGCTGTACCCCTGGACAACCAGCCCTCCATCTCTTGAGATCTAGTCAGAGCCTTATATA  
T Q S P L S L P V T L G Q P A S I S C R S S Q S L I Y

779 TAGTGATGGAAACGCCCTATTACATTGGTTCTTGAGAGCCAGGCCAGTCTCCAGGCTCCTGATCTACAAAGTTTCCA  
S D G N A Y L H W F L Q K P G Q S P R L L I Y K V S N

859 ACCGATTTTCTGGGGTCCAGACAGGTTCACTGGCAGTGGATCTGGACAGATTTCACACTCAAGATCAGCAGAGTGGAG  
R F S G V P D R F S G S G S G T D F T L K I S R V E

939 GCCGAGGATGTGGGAGTTTATTACTGTTCTCAAAGTACACATGTTCCGTGGACGTTCCGTCAGGCACCAAGGTGGAAAT  
A E D V G V Y Y C S Q S T H V P W T F G Q G T K V E I

1019 CAAACGTGAGTAGAATTTAATCTAGAAATTTCTAACTCTGAGGGGGTGGGATGACGTGGCCATTCTTTGGCTAAAGCAAT  
K R

1099 GAGTTTACTGCAAGGTCAGAAAAGCATGCAAAGCCCTCAGAAATGGCTGCAAAGAGCTCCAAACAAAACAAATTTAGAACTTT  
1179 ATTAAGGAATAGGGGGAAGCTAGGAAGAAATCAAACATCAAGATTTTAAATACGCTTCTTGCTCTCTTGCTATAATT  
1259 ATCTGGGATAAGCATGCTGTTTTCTGCTGCTCCCTAACATGCCCTGTGATTATCCGCAAAACAACACCCCAAGGGCAGAA  
1339 CTTTGTACTTAAACACCATCCTGTTTGTCTTCTTCTCTCAGGAACGTGGCTGCACCATCTGTCTTCACTTCCCGCCAT  
T V A A P S V F I P P S

1419 CTGATGAGCAGTTGAAATCTGGAACTGCCCTCTGTTGTGTGCTGCTGAATAACTTCTATCCAGAGAGGCCAAAGTACAG  
D E Q L K S G T A S V V C L L N N F Y P R E A K V Q W

1499 TGGAAAGGTGGATAACGCCCTCCAATCGGGTAATCTCCAGGAGAGTGTCAAGAGCAGGACAGCAAGGACAGCACTACAG  
K V D N A L Q S G N S Q B S V T E Q D S K D S T Y S L

1579 CCTCAGCAGCACCCTGACGCTGAGCAAGCAGACTACGAGAAACAAAGTCTACGCTGCGAAGTCACCCATCAGGGCC  
S S T L T L S K A D Y E K H K V Y A C E V T H Q G L

1659 TGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTAGAGGGAGAAATGCCCCACCTGCTCCTCAGTTCCAGC  
S S P V T K S F N R G B C

1739 CTGACCCCTCCCATCTCTTGGCTCTGACCCCTTTTCCACAGGGGACCTACCCCTATTGCGGTCTCCAGCTCATCTTT  
1819 CACCTCACCCCTCTCTCTCTCTGCTTAAATATGCTAATGTTGGAGGAGAAATGAATAAATAAGTGAATCTTTGACAC  
1899 CTGTGGTTTCTCTCTTCTCTCTTAAATATATATCTGTTGTTTACCAACTACTCAATTTCTCTTATAGGGGACTAA  
1979 ATATGATGTCATCTAAGGCGCATAAACATTTATAAAATCATCTTCTATTTTACCTATCATCTCTGCAAGAC  
2059 AGTCTCTCCCTCAAACCCACAAGCCTTCTGCTCTCAAGTCCCCTGGGCCATGTAGGAGAGACTTGCTTCTGTTTTC  
2139 CCTCCTCAGCAAGCCCTCATAGTCTCTTTTAAAGGTGACAGGCTTACAGTCATATATCTTTGATTCAATTCCTGAGA  
2219 ATCAACCAAGCAAAATTTTCAAAGAGAAACCTGCTATAAGAGAATCATTCATTGCAACATGATATAAATAACAAC  
2299 ACAATAAAGCAATTAATAAACAACATAGGGAAATGTTTAAAGTTCATCATGGTACTTAGACTTAATGGAATGTCATG  
2379 CCTTATTTACATTTTAAACAGGTACTGAGGGACTCTGCTGCGCAAGGGCCGATTAGTACTTTCCACAACCTAATTT  
2459 AATCCACACTATATCTGTGAGATTAAAAACATTCATTAAATGTTGCAAGGTTCTATAAGCTGAGAGACAAATATATTC

2539 TATAACTCAGCAATCCCACTTCTAGGATCC

The complete sequence of the Hu266 light chain gene is located between the MluI and BamHI sites in pVk-Hu266. The nucleotide number indicates its position in pVgk-Hu266. The V<sub>k</sub> and C<sub>k</sub> exons are translated in single letter code; the dot indicates the translation termination codon. The mature heavy chain starts at the double-underlined aspartic acid (D). The intron sequences are in italic.

Figure 5. pVg1-Hu266 polynucleotide sequences for expressing humanized 266 heavy chain and single amino acid codes for expressed humanized 266 heavy chain

```

619 ACGCGTCCACCATGAATTCGGGCTCAGCTTGATTTCTTGTCCTTGTTTAAAGGTGCTCTGTGTGAAGTGCAGCTG
    M N F G L S L I F L V L V L K G V L C E V Q L
699 GTGGAGTCTGGGGGAGGTTTAGTGCAGCCTGGAGGCTCCCTGAGACTCTCCTGTGCAGCCTCTGGATTCACTTTTAGTAG
    V E S G G G L V Q P G G S L R L S C A A S G F T F S R
779 GTATTCCATGCTCTGGGTTGCCAGGCTCCAGGCAAGGGCCTGGAATTGGTGCACAAATTAATAGTGTGGTAATAGCA
    Y S M S W V R Q A P G K G L E L V A Q I N S V G N S
859 CCTACTATCCAGACACTGTAAAGGCGGATTACCATCTCCAGAGCAATGCCAAGAACCCCTGTACCTGCAAAATGAAC
    T Y Y P D T V K G R P T I S R D N A K N T L Y L Q M N
939 TCCCTGAGGGCCGAAGACACGGCGCTGTATTACTGTGCAAGCGGAGACTACTGGGGCCAAGGCACCTCTGGTGACAGTCTC
    S L R A E D T A V Y Y C A S G D Y W G Q G T L V T V S
1019 CTCAGGTGAGTCTCTCAACCTCTAGAGCTTTCTGGGGCAGGCCAGGCCCTGACCTTGGCTTTGGGGCAGGGAGGGGGCTA
    S
1099 AGGTGAGGCAGGTGGCGCCAGCCAGGTGCACCCAAATGCCCATGAGCCGAGACACTGGAAGCTGAACTCTCGCGACAGT
1179 TAAGAAACCCAGGGGCTCTGCGCCCTGGGCCCAGCTCTGTCCCAACCGCGGTACATGGCAACCACTCTCTTGCACTCT
    A
1259 CCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCTCTCCAGAGCACCTCTGCGGGCACAGCGGCCCTGGGCTGCCTG
    S T K G P S V F P L A P S S K S T S G G T A A L G C L
1339 GTCAAGGACTACTTCCCGAACCGGTGACGGTGTCTGGAACTCAGGCGCCCTGACAGCGGGGTGCACACCTTCCCGGC
    V K D Y F P E P V T V S W N S G A L T S G V H T F P A
1419 TGTCTACAGTCTCTCAGGACTCTACTCCCTCAGCAGCGTGGTACCGTCCCTCCAGCAGCTTGGGCAACCCAGACCTACA
    V L Q S S G L Y S L S S V V T V P S S S L G T Q T Y
1499 TCTGCAACGTGAATCACAAGCCAGCAACCAAGGTGGACAGAAAGTTGGTGAGAGGCCAGCACAGGGAGGGAGGGGTG
    I C N V N H K P S N T K V D K K V
1579 TCTGCTGGAAGCCAGGCTCAGCGCTCTGCTGGACGCAATCCCGGCTATGCAAGCCCACTCCAGGGGAGCAAGGCAGGCC
1659 CCGTCTGCTCTTCAACCGGAGGCTCTGCGCGCCCACTCATGCTCAGGGAGAGGGTCTTCTGGCTTTTCCCGAGGCT
1739 CTGGGCGAGGACAGGCTAGGTGCGCCCTAACCCAGGCGCTGCACAAAGGGGCAAGGTCTCGGCTCAGACCTGCCAAGAG
1819 CCATATCGGGGAGGACCTGCGCCCTGACCTAAGCCACCCCAAGGCCAACTCTCCAATCCCTCAGCTCGGACACCTTC
1899 TCTCTCCAGATTCCAGTAACCTCCAATCTTCTCTCTGCAAGCCCAATCTTGTGACAAACTCACAATGCCACCG
    E P K S C D K T H T C P P
1979 TGCCAGGTAAAGCCAGCCAGGCTCTGCGCTCCAGCTCAAGGCGGGAAGGTGCGCTAGAGTAGCCTGCATCCAGGGACA
    C P
2059 GGCCCCAGCGGGTGTGACAGCTCCACCTCCATCTCTTCTCAGCACTGAACCTCTGGGGGACCGTCACTCTTCTC
    A P E L L G O P S V F L
2139 TTCCCCCAAAACCAAGGACACCTCATGATCTCCCGGACCCCTGAGGTCACTGCTGTGTGTGGACGTGAGCCAGCA
    F P P K P K D T L M I S R T P E V T C V V V D V S E E
2219 AGACCTGAGGTCAAGTCAACTGGTACGTGGAGCGGTGGAGGTGCATAATGCCAAGACAAGCCGCGGGAGGAGCAGT
    D P E V K F N W Y V D G V E V H N A K T K P R E E Q
2299 ACAACAGCACGTACCGTGTGTGAGCGTCTCACCGTCTGCAACAGGACTGGCTGAATGGCAAGGAGTACAAGTCAAG
    Y N S T Y R V V S V L T V L H Q D W L N G K E Y K C K
2379 GTCTCCAAACAAGCCCTCCAGCCCCCATCGAGAAACCATCTCCAAAGCCAAAGGTGGGACCCGTGGGTGGAGGGGCC
    V S N K A L P A P I E K T I S K A K
2459 ACATGGACAGAGGCGCGCTCGGCCACCCCTCTGCGCTGAGAGTGACCGCTGTACCAACCTCTGTCCCTACAGGGCAGCCC
    G Q P
2539 CGAGAACCACAGGTGTACACCTTGCCCCCATCCCGGATGAGCTGACCAAGAACAGGTGAGCTGACCTGCTGCTGCA
    R E P Q V Y T L P P S R D E L T K N Q V S L T C L V K
2619 AGGCTTCTATCCAGCGACATCGCGTGGAGTGGAGAGCAATGGGCGAGCCGAGAACAACTACAAGACCAGCGCTCCCG

```

G P Y P S D I A V E W E S N G Q P E N N Y K T T P P  
 2699 TGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTC  
 V L D S D G S F F L Y S K L T V D K S R W Q Q G N V F  
 2779 TCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGAGTGG  
 S C S V N H B A L H N H Y T Q K S L S L S P G K  
 2859 ACGGCGGCAAGCCCCGCTCCCGGGCTCTCGCGTCCGACAGGATGCTTGGCAGTACCCCCGTACATACTTCCC  
 2939 GCGGCCAGCATGGAATAAAGCACCCAGCGCTGCTTGGGCCCCCTGCGAGACTGTGTGTCTTTCACGGGTCTGSC  
 3019 CGAGTCTGAGGCCCTGAGTGGCATGAGGGAGGCAGAGCGGGTCCCCTGCTCCCCCACTGGCCAGGCTGTGCAAGTGTG  
 3099 CTGGGCGGCTAGGGTGGGGCTCAGCCAGGGCTGCCCTCGGAGGGTGGGGATTGCCAGCGTGGCCCTCCCTCCAGC  
 3179 AGCACCTGCCCTGGGCTGGGCCACGGGAAGCCCTAGGAGCCCTGGGGACAGACACACAGCCCTGCTCTGTAGGAGC  
 3259 TGTCTGTCTCTGTAGCCGCCCTGTCTCCGACCTCCATGCCCTAGTGGGGCATGCTAGTCCATGTGGTAGGACAGG  
 3339 CCTCCCTCACCCTCTACCCCCACGGCACTAACCCCTGGCTGCCCTGCCAGCCTCGCACCCGCTGGGGACACAAACG  
 3419 ACTCCGGGACATGCACTCTCGGGCCCTGTGGAGGAGTGGTGAGATGCCACACACACTCAGCCAGACCCGTTCA  
 3499 ACAAAACCCGCACTGAGGTGGCCGGCCACAGGCCACACACACACACGTCACGCTCAGACAGGAGCTCAGCCGG  
 3579 GCGAACTGCACAGCACCCAGACCAGGCAAGGTCTCTGCACACGTGAACACTCTCGGACACAGGCCCCAGGAGCCCCA  
 3659 CGGGCACCTCAAGGCCACGAGCCTCTCGGCAGCTTCTCCATGCTGACCTGCTCAGACAAACCCAGCCCTCTCTCA  
 3739 CAAGGGTGCCCTGACGCGGCCACACACACAGGGGATCACACCCAGCTCACGTCCCTGGCCCTGGGCCACTTCCAG  
 3819 TGCCGCCCTTCCCTGCAGGATCC

The complete sequence of the Hu266 heavy chain gene is located between the MluI and BamHI sites in pVg1-Hu266. The nucleotide number indicates its position in pVg1-Hu266. The  $V_H$  and  $C_H$  exons are translated in single letter code. The dot indicates the translation termination codon. The mature heavy chain starts at the double-underlined glutamic acid (E). The intron sequences are in italics.

Figure 6. Plasmid pVkJHu266






Figure 7 Plasmid pVg1-Hu266





Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) EP 1 481 992 A3

(12) EUROPEAN PATENT APPLICATION

(88) Date of publication A3:  
08.12.2004 Bulletin 2004/50

(43) Date of publication A2:  
01.12.2004 Bulletin 2004/49

(21) Application number: 04011466.2

(22) Date of filing: 26.02.2001

(51) Int Cl.7: C07K 16/18, C12N 15/13,  
C12N 5/10; A61K 39/00,  
A61P 25/28

(84) Designated Contracting States:  
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE TR  
Designated Extension States:  
AL LT LV MK RO SI

(30) Priority: 24.02.2000 US 184601 P  
08.12.2000 US 254465 P  
08.12.2000 US 254498 P

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
01913081.4 / 1 257 584

(71) Applicants:  
• WASHINGTON UNIVERSITY ST. LOUIS  
St. Louis Missouri 63110 (US)  
• Eli Lilly & Company  
Indianapolis, IN 46285 (US)

(72) Inventors:  
• Holtzman, David, M.  
St. Louis MO 63141 (US)  
• Demattos, Ronald  
St. Louis MO 63110 (US)  
• Bales, Kelly, R.  
Cloverdale IN 46120 (US)  
• Paul, Steven, M.  
Carmel IN 46032 (US)  
• Tsurushita, Naoya  
Palo Alto CA 94306 (US)  
• Vasquez, Maximiliano  
Palo Alto CA 94303 (US)

(74) Representative: Grund, Martin, Dr. et al  
Dr. Volker Vossius,  
Patentanwaltskanzlei,  
Geibelstrasse 6  
81679 München (DE)

(54) Humanized antibodies that sequester amyloid beta peptide

(57) A method to treat conditions characterized by formation of amyloid plaques both prophylactically and therapeutically is described. The method employs humanized antibodies which sequester soluble A $\beta$  peptide from human biological fluids or which preferably specifically bind an epitope contained within position 13-28 of the amyloid beta peptide A $\beta$ .




Figure 1

EP 1 481 992 A3



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 04 01 1466

## DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Y	WINTER ET AL: "HUMANIZED ANTIBODIES" IMMUNOLOGY TODAY, ELSEVIER PUBLICATIONS, CAMBRIDGE, GB, vol. 14, no. 6, 1996, pages 243-246, XP001005438 ISSN: 0167-4919 * See pages 243-245 (Building humanized antibodies) *	1-13	C07K16/18 C12N15/13 C12N5/10 A61K39/00 A61P25/28
Y	SEUBERT ET AL: "ISOLATION AND QUANTIFICATION OF SOLUBLE ALZHEIMER'S BETA-PEPTIDE FROM BIOLOGICAL FLUIDS" NATURE, MACMILLAN JOURNALS LTD. LONDON, GB, vol. 359, no. 6393, 24 September 1992 (1992-09-24), pages 325-327, XP000616173 ISSN: 0028-0836 * See the antibody 266 *	1-13	
Y	PILLOT ET AL: "Fusogenic properties of the C-terminal domain of the Alzheimer beta-amyloid peptide" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 271, 1996, pages 28757-28765, XP002299386 * See references to the central domain (13-28) throughout the article *	1-13	TECHNICAL FIELDS SEARCHED (Int.Cl.7) C07K C12N A61K A61P
Y	ST GEORGE-HYSLOP ET AL: "Antibody clears senile plaque" NATURE, vol. 400, July 1999 (1999-07), pages 116-117, XP002299786 * See page 116 (clearance of amyloid-beta peptide) *	1-13	
The present search report has been drawn up for all claims			
Place of search Berlin		Date of completion of the search 8 October 2004	Examiner Korsner, S-E
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document			

EPO FORM 1502 (01.02.2004)





European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 04 01 1466

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
P, Y, D	<p>BARD ET AL: "Peripherally administered antibodies against amyloid beta-peptide enter the central nervous system and reduce pathology in a mouse model of Alzheimer disease". NATURE MEDICINE, NATURE PUBLISHING, CO, US, vol. 6, no. 8, August 2000 (2000-08), pages 916-919, XP002154518 ISSN: 1078-8956 * See the Summary *</p> <p>-----</p>	1-13	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
The present search report has been drawn up for all claims			
Place of search Berlin		Date of completion of the search 8 October 2004	Examiner Korsner, S-E
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p>		<p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>..... &amp; : member of the same patent family, corresponding document</p>	

EPO FORM 1503 03/02 (PMD01)